

DESCRIPTION

DIFLUOROMETHYLTRIAZOLONE COMPOUNDS, USE OF THE SAME AND INTERMEDIATES FOR THE PRODUCTION THEREOF

5

Technical Field

The present invention relates to triazolone compounds, use thereof and production intermediates thereof.


10 Background Art

WO99/07687 describes that a certain kind of triazolone compounds has a fungicidal activity.

Disclosure of the Invention

15 The present invention provides compounds having a fungicidal activity, particularly an excellent fungicidal activity on fungi causing plant diseases and useful for controlling plant diseases.

Namely, the present invention provides a triazolone
20 compound of the formula [I] (hereinafter, referred to as the compound of the invention):



wherein,

R^1 represents a A^1-L^1- , $A^1-ON=CA^2-$, $A^1-ON=C(Me)CH_2ON=CA^2-$,
25 $A^1-C(A^2)=N-OCH_2-$, $A^1S-C(A^2)=N-$, $A^1-C(=S)NH-$, $A^1S-C(=S)NH-$,
 $A^1S-C(SA^2)=N-$, $A^1-ON=C(CN)-$, $A^1-ON=C(Me)CH_2ON=C(CN)-$,
 $A^1-C(CN)=N-OCH_2-$, halogen atom, nitro or cyano;

wherein, L^1 represents single bond, oxygen atom, sulfur atom, carbonyl, $-OCH_2-$, $-SCH_2-$, $-C(=O)O-$, $-OC(=O)-$, $-C(=O)OCH_2-$, $-NH-$ or C_1-C_6 alkylimino;

A^1 and A^2 , which are the same or different, represent
5 hydrogen atom, C_1-C_{10} alkyl, C_2-C_{10} alkenyl, C_2-C_{10} alkynyl, C_3-C_{10} cycloalkyl, (C_3-C_{10} cycloalkyl) C_1-C_{10} alkyl, C_5-C_{10} cycloalkenyl, (C_5-C_{10} cycloalkenyl) C_1-C_{10} alkyl, phenyl, naphthyl, phenyl C_1-C_{10} alkyl, naphthyl C_1-C_{10} alkyl, 5- or 6-membered heterocyclic group optionally condensed with
10 a benzene ring, or methyl substituted by 5- or 6-membered heterocyclic group optionally condensed with a benzene ring;

the alkyl, the alkenyl, the alkynyl, the cycloalkyl, the cycloalkylalkyl, the cycloalkenyl and the
15 cycloalkenylalkyl, represented by A^1 and A^2 , may optionally be each substituted by one or more substituents selected from the group consisting of halogen atom(s), cyano, nitro, C_1-C_{10} alkoxy, C_1-C_{10} haloalkoxy, C_1-C_{10} alkylthio, C_1-C_{10} haloalkylthio, (C_1-C_9 alkyl) carbonyl, (C_1-C_9 alkoxy) carbonyl, (C_1-C_9 alkyl) carbonylamino, phenyl, phenoxy, benzyloxy and tri(C_1-C_{10} alkyl) silyl;

the phenyl, the naphthyl, the benzene ring in the phenylalkyl, the naphthalene ring in the phenylnaphthyl, the heterocyclic group, and the heterocyclic ring in the
25 methyl substituted by heterocyclic group, represented by A^1 and A^2 , may optionally be each substituted by one or more substituents selected from the group consisting of halogen atom(s), cyano, nitro, C_1-C_{10} alkyl, C_1-C_{10} haloalkyl, C_3-C_{10} cycloalkyl, C_1-C_{10} alkoxy, C_1-C_{10} haloalkoxy, C_1-C_{10} alkylthio, C_1-C_{10} haloalkylthio, (C_1-C_9
30


alkyl) carbonyl, (C₁-C₉ alkoxy) carbonyl, (C₁-C₉ alkyl) carbonylamino, phenyl, phenoxy, benzyloxy, tri(C₁-C₁₀ alkyl) silyl, methylenedioxy and difluoromethylenedioxy; with the proviso, when L¹ is single bond, A¹ is not a hydrogen atom;

T represents optionally substituted *m*-phenylene, optionally substituted *m*-azaphenylene (*m*-pyridinediyl) or optionally substituted *m*-diazaphenylene (*m*-diazinediyl, namely, *m*-pyridazinediyl, *m*-pyrimidinediyl or *m*-pyrazinediyl) bonded to R¹ and to CH₂ each via a carbon atom;


wherein the substituents are one or more substituents selected from the group consisting of halogen atoms, cyano, nitro, C₁-C₆ alkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, C₁-C₆ alkylthio, C₁-C₆ haloalkylthio and (C₁-C₅ alkoxy) carbonyl;

and a fungicidal composition containing this compound as an active ingredient.


The present invention provides also 5-difluoromethyl-2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one of the formula [L]:




; a difluoroacetyl semicarbazide compound of the formula [XXVI]:




wherein R¹ and T have the same meanings as described above; a semicarbazide compound of the formula [VII]:



wherein R^1 and T have the same meanings as described above;
2-methyl-5-phenylbenzylamine of the formula [XVI-1]:




5 , inorganic acid salt (for example, hydrochloric acid addition salt, hydrobromic acid addition salt, sulfuric acid addition salt) and sulfonic acid salt (for example, methanesulfonic acid addition salt) of 2-methyl-5-phenylbenzylamine;
2-methyl-5-phenylbenzyl chloride of the formula [XIII-1]:



10


, a sulfonate compounds of the formula [XIII-2]:



wherein R^9 represents a methyl or p-tolyl;

N,N-dimethyl-(2-methyl-5-phenylbenzyl)amine of the formula

15 [LI]:



, inorganic acid salt (for example, hydrochloric acid addition salt, hydrobromic acid addition salt, sulfuric acid addition salt) and sulfonic acid salt (for example, methanesulfonic acid addition salt) of N,N-dimethyl-(2-methyl-5-phenylbenzyl)-
5 amine, useful as a production intermediate for a compound of the invention.

In the present invention;
as the C₁-C₁₀ alkyl represented by A¹ and A², for example, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, 1-methylpropyl,
10 pentyl, 1-methylbutyl, 1-ethylbutyl, 2-methylbutyl, 3-methylbutyl, 2,2-dimethylpropyl, 1,2-dimethylpropyl, 1,1-dimethylpropyl, hexyl, 1-methylpentyl, 1-ethylpentyl, 3,3-dimethylbutyl, heptyl, 3,7-dimethyloctyl and the like are listed; as the C₂-C₁₀ alkenyl represented by A¹ and A², for example,
15 vinyl, allyl, 1-methyl-2-propenyl, 2-methyl-2-propenyl, 2-butenyl, 2-pentenyl, 3-methyl-2-butenyl and the like are listed; as the C₂-C₁₀ alkynyl represented by A¹ and A², for example, ethynyl, propargyl, 1-methyl-2-propynyl, 2-butyne and the like are listed; as the C₃-C₁₀ cycloalkyl represented by A¹ and A²,
20 for example, cyclopropyl, cyclopentyl, cyclohexyl and the like are listed; as the (C₃-C₁₀ cycloalkyl) alkyl represented by A¹ and A², for example, cyclopropylmethyl, cyclopentylmethyl, 2-cyclopentylethyl, cyclohexylmethyl and the like are listed; as the C₅-C₁₀ cycloalkenyl represented by A¹ and A², for example,
25 cyclopentenyl, cyclohexenyl and the like are listed; as the (C₅-C₁₀ cycloalkenyl) alkyl represented by A¹ and A², for example, cyclopenten-1-ylmethyl, cyclohexen-1-ylmethyl and the like are listed; as the phenyl C₁-C₁₀ alkyl represented by A¹ and A², for example, phenylmethyl, 2-phenylethyl, 3-phenylpropyl,
30 4-phenylbutyl and the like are listed; as the naphthyl C₁-C₁₀

alkyl represented by A¹ and A², for example, α-naphthylmethyl, β-naphthylmethyl and the like are listed; as the 5- or 6-membered heterocyclic group optionally condensed with a benzene ring represented by A¹ and A², for example, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 2-pyrazinyl, 3-pyridazinyl, 4-pyridazinyl, 2-thienyl, 3-thienyl, 2-furyl, 3-furyl, pyrrol-1-yl, pyrrol-2-yl, pyrrol-3-yl, 1-pyrazolyl, 3-pyrazolyl, 4-pyrazolyl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 3-isothiazolyl, 4-isothiazolyl, 5-isothiazolyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, 3-isooxazolyl, 4-isooxazolyl, 5-isooxazolyl, 1-imidazolyl, 2-imidazolyl, 4-imidazolyl, 5-imidazolyl, 1-(1,2,4-triazolyl), 3-(1,2,4-triazolyl), 4-(1,2,4-triazolyl), 2-benzothienyl, 3-benzothienyl, benzothiazol-2-yl, 2-quinolyl and the like are listed; and as the methyl substituted by a 5- or 6-membered heterocyclic group optionally condensed with a benzene ring represented by A¹ and A², for example, 2-pyridylmethyl, 4-pyridylmethyl, 2-pyrimidinylmethyl, 4-pyrimidinylmethyl, 3-pyrazolylmethyl, 2-thiazolylmethyl, 2-imidazolylmethyl, 3-(1,2,4-triazolyl)methyl, 2-quinolylmethyl and the like are listed.

Here, the alkyl, the alkenyl, the alkynyl, the cycloalkyl, the cycloalkylalkyl, the cycloalkenyl and the cycloalkenylalkyl represented by A¹ and A² may optionally be each substituted by one or more substituents selected from the group consisting of halogen atom(s) (fluorine atom, chlorine atom, bromine atom, iodine atom), cyano, nitro, C₁-C₁₀ alkoxy, C₁-C₁₀ haloalkoxy, C₁-C₁₀ alkylthio, C₁-C₁₀ haloalkylthio, (C₁-C₉ alkyl) carbonyl, (C₁-C₉ alkoxy) carbonyl, (C₁-C₉ alkyl) carbonylamino, phenyl, phenoxy, benzyloxy and tri(C₁-C₁₀ alkyl) silyl; and as the alkoxy,

for example, a methoxy, ethoxy, propoxy, isopropoxy, butoxy, sec-butoxy, isobutoxy, pentyloxy and the like are listed; as the haloalkoxy, for example, a trifluoromethoxy, difluoromethoxy, bromodifluoromethoxy, chlorodifluoromethoxy, fluoromethoxy, 2,2,2-trifluoroethoxy, 1,1,2,2-tetrafluoroethoxy and the like are listed; as the alkylthio, for example, a methylthio, ethylthio, propylthio, butylthio, isobutylthio, sec-butylthio, pentylthio, hexylthio and the like are listed, as the haloalkylthio, for example, a trifluoromethylthio, difluoromethylthio, bromodifluoromethylthio, chlorodifluoromethylthio, fluoromethylthio, 2,2,2-trifluoroethylthio, 1,1,2,2-tetrafluoroethylthio and the like are listed; as the alkylcarbonyl, for example, an acetyl, propanoyl, butanoyl, 3-methylbutanoyl and the like are listed; as the alkoxycarbonyl, for example, a methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, sec-butoxycarbonyl, pentyloxycarbonyl, hexyloxycarbonyl and the like are listed; as the alkylcarbonylamino, for example, an acetylamino, propanylamino, butanoylamino, 3-methylbutanoylamino and the like; and as the C₃-C₂₀ trialkylsilyl, for example, a trimethylsilyl, triethylsilyl and the like are listed.


Further, the phenyl, the naphthyl, the benzene ring in the phenylalkyl, the naphthalene ring in the phenylnaphthyl, the heterocyclic group, and the heterocyclic ring in the methyl substituted by a heterocyclic group, represented by A¹ and A², may optionally be each substituted by one or more substituents selected from the group consisting of halogen atom(s) (fluorine atom, chlorine atom, bromine atom, iodine atom), cyano, nitro,

C₁-C₁₀ alkyl, C₁-C₁₀ haloalkyl, C₃-C₁₀ cycloalkyl, C₁-C₁₀ alkoxy, C₁-C₁₀ haloalkoxy, C₁-C₁₀ alkylthio, C₁-C₁₀ haloalkylthio, (C₁-C₉ alkyl) carbonyl, (C₁-C₉ alkoxy) carbonyl, (C₁-C₉ alkyl) carbonylamino, phenyl, phenoxy, benzyloxy, tri(C₁-C₁₀ alkyl) silyl, methylenedioxy and difluoromethylenedioxy; and as the alkoxy, the haloalkoxy, the alkylthio, the haloalkylthio, the alkylcarbonyl, the alkoxy carbonyl, the alkylcarbonylamino and the trialkylsilyl, the same substituents as described above are listed; as the alkyl, for example, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, 1-methylpropyl, pentyl, 1-methylbutyl, 1-ethylbutyl, 2-methylbutyl, 3-methylbutyl, 2,2-dimethylpropyl, 1,2-dimethylpropyl, 1,1-dimethylpropyl, hexyl, 1-methylpentyl, 1-ethylpentyl, 3,3-dimethylbutyl, heptyl, 3,7-dimethyloctyl and the like are listed; as the haloalkyl, for example, trifluoromethyl, 2,2,2-trifluoroethyl, 1,1,2,2-tetrafluoroethyl and the like are listed; and as the cycloalkyl, for example, cyclopropyl, cyclopentyl, cyclohexyl and the like are listed; and the methylenedioxy and the difluoromethylenedioxy are each substituted on adjacent two carbon atoms in a benzene ring, naphthalene ring or heterocyclic ring.

The *m*-phenylene, *m*-azaphenylene and *m*-diazaphenylene, bonded to R¹ and to CH₂ each via a carbon atom, represented by T may optionally be substituted by halogen atoms (fluorine atom, chlorine atom, bromine atom, iodine atom), cyano, nitro, C₁-C₆ alkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ haloalkoxy, C₁-C₆ alkylthio, C₁-C₆ haloalkylthio or (C₁-C₅ alkoxy) carbonyl; and as the alkyl, for example, methyl, ethyl and the like are listed; as the haloalkyl, for example, fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl and the like are listed; as the

alkoxy, for example, methoxy, ethoxy and the like are listed; as the haloalkoxy, for example, trifluoromethoxy, difluoromethoxy, bromodifluoromethoxy, chlorodifluoromethoxy, fluoromethoxy, 2,2,2-trifluoroethoxy, 1,1,2,2-tetrafluoroethoxy and the like are listed; as the alkylthio, for example, methylthio, ethylthio and the like are listed; as the haloalkylthio, for example, a trifluoromethylthio, difluoromethylthio, bromodifluoromethylthio, chlorodifluoromethylthio, fluoromethylthio, 2,2,2-trifluoroethylthio, 1,1,2,2-tetrafluoroethylthio and the like are listed; and as the alkoxycarbonyl, for example, methoxycarbonyl, ethoxycarbonyl and the like are listed.

As more specific examples of T, groups of the following formulae:




are listed.


Among the compounds of the invention, the compounds in which R¹ is optionally substituted phenyl (namely, R¹ is a A¹-L¹-, L¹ is single bond, and A¹ is optionally substituted phenyl) are preferable because of an excellent fungicidal effect. Further, the compounds in which T is *m*-phenylene or substituted

m-phenylene are preferable because of an excellent fungicidal effect, and the preferable substituents include methyl, fluorine atom, chlorine atom and trifluoromethyl.

Particularly, the compounds of the following formula
 5 [LX] in which R¹ is optionally substituted phenyl and T is methylphenylene described below are preferable because of an excellent fungicidal effect.



As the specific example of such compounds,
 10 5-difluoromethyl-2-methyl-4-(2-methyl-5-phenylbenzyl)-2,4-dihydro-3H-1,2,4-triazol-3-one [Compound 1] of the formula:




is mentioned.

15 The compound of the invention can be produced according to the following Production Method A to Production Method G. Depending on the definition of R¹, a desired R¹ can be introduced or structured last (for example, Production Method H). In these Production Methods, a protective group can be used for protecting
 20 a functional group from reaction, if necessary.

Production Method A

The compound of the invention of the formula [I] can be produced from a formyl compound of the formula [II] by, for example, a method shown in the following chemical reaction formula:




wherein, R^1 and T have the same meanings as described above.

The reaction temperature of the reaction of obtaining the compound of the invention of the formula [I] from the formyl compound of the formula [II] is usually in the range from -20°C to 100°C , and the reaction time is usually in the range from 10 minutes to 10 hours. The ratio of diethylaminosulfur trifluoride used for this reaction is usually 1 to 20 mol per mol of the formyl compound of the formula [II].

The reaction is conducted usually in a solvent, and as the solvent used, for example, halogenated hydrocarbons such as chloroform, chlorobenzene and the like, ethers such as 1,4-dioxane, tetrahydrofuran, diethyl ether and the like, aliphatic hydrocarbons such as hexane, petroleum ether and the like, aromatic hydrocarbons such as toluene, xylene and the like, and mixtures thereof are listed.

The reaction solution after completion of the reaction is subjected to post treatments such as, for example, washing with a sodium bicarbonate solution before concentration of the organic layer, and the like, giving isolation of the intended compound. This intended compound can also be purified by re-crystallization, chromatography and the like.

The formyl compound of the formula [II] can be produced, for example, according to the following scheme.



wherein, R^1 and T have the same meanings as described above,
 5 L^2 represents a leaving group such as a chlorine atom, bromine atom, iodine atom, p-toluenesulfonyloxy, methanesulfonyloxy, trifluoromethanesulfonyloxy and the like, and R^5 represents a C_1 - C_5 alkyl (for example, methyl, ethyl and the like), phenyl or trihaloethyl (for example, trichloroethyl and the like).

10 A semicarbazide compound of the formula [VII] can be produced by reacting an isocyanate compound of the formula [III] with methylhydrazine. The reaction temperature of the reaction

is usually in the range from -20°C to 50°C, and the reaction time is usually in the range from 1 to 100 hours. The ratio of methylhydrazine used for this reaction is usually 1 to 10 mol per mol of the isocyanate compound of the formula [III].

5 The reaction is usually conducted in a solvent, and as the solvent used, for example, ethers such as 1,4-dioxane, tetrahydrofuran, diethyl ether and the like, halogenated hydrocarbons such as chloroform, chlorobenzene and the like, aliphatic hydrocarbons such as hexane, petroleum ether and the like, aromatic
10 hydrocarbons such as toluene, xylene and the like, amides such as N,N-dimethylformamide and the like, dimethylsulfoxide and the like, and mixtures thereof are listed. The reaction solution after completion of the reaction is subjected to post treatments such as concentration, and the like, giving isolation
15 of the intended compound. This intended compound can also be purified by re-crystallization, chromatography and the like. A semicarbazide compound of the formula [VIII] can be produced by reacting the isocyanate compound of the formula [III] with 1-methyl-2-hydroxyacetylhydrazine in the same manner as
20 described above. A semicarbazide compound of the formula [IV] can also be produced by reacting the isocyanate compound of the formula [III] with hydroxyacetylhydrazine in the same manner as described above.

A compound of the formula [V] can be obtained by reacting
25 the semicarbazide compound of the formula [IV] with a base, and for example, can be obtained by treating the compound of the formula [V] with an aqueous potassium hydroxide solution.


A compound of the formula [VI] can be obtained by reacting the semicarbazide compound of the formula [VIII] with a base,
30 and for example, can be obtained by treating the compound of

the formula [VI] with an aqueous potassium hydroxide solution.

A compound of the formula [VI] can be obtained by reacting the compound of the formula [V] with a base, and as the base used, for example, potassium carbonate is mentioned.

5 The compound of the formula [II] can be obtained by oxidizing the compound of the formula [VI], and as the oxidizing agent used, for example, manganese dioxide is mentioned.

The isocyanate compound of the formula [III] can be produced, for example, according to the following scheme.



10

wherein, R^1 , L^2 and T have the same meanings as described above, R^6 represents a C_1 - C_4 alkyl (for example, methyl, ethyl and the like), M^1 represents silver or sodium, DIBAL represents diisobutyl aluminum hydride, BuLi represents butyllithium and DMF represents N,N-dimethylformamide.

15

The reaction of reacting an amine compound of the formula [XVI] with triphosgene, diphosgene or phosgene to obtain the isocyanate compound of the formula [III] can be conducted, for example, by a method described in J. Org. Chem. 61, 3883-3884 (1996) or by a method described in Intermediate Production Example 3 described later.

The reaction of reducing a nitrile compound of the formula [XV] to obtain the amine compound of the formula [XVI] can be conducted, for example, by a method of reduction using aluminum lithium hydride (LiAlH_4), by a method of hydrogenating the nitrile compound of the formula [XV] in the presence of a catalyst, and by the like manner; as the catalyst used in hydrogenation, for example, Raney nickel, palladium and the like are listed. Regarding the method of using Raney nickel, JP-A No. 8-291116 can be referred, and regarding the method of using palladium, J. Am. Chem. Soc., 50, 3370 (1928) can be referred, respectively.

The reaction of obtaining a compound of the formula [XIII] wherein L^2 represents a p-toluenesulfonyloxy, methanesulfonyloxy or trifluoromethanesulfonyloxy from an alcohol compound of the formula [XII] can be conducted, for example, according to a method described in Intermediate Production Example 12 described later.


The reaction of obtaining a compound of the formula [XIII] wherein L^2 represents a chlorine atom from the alcohol compound of the formula [XII] can be conducted, for example, by a method of reacting $\text{CCl}_4\text{-PPh}_3$ to the alcohol compound of the formula [XII] (specifically, method described in Intermediate Production Example 9 described later), by a method of reacting conc. Hydrochloric acid to the alcohol compound of the formula [XII] (see, Org. Synth., IV, 576 (1967)), and by the like manner.

The similar methods are applicable to the compound which L2 is bromine atom or iodine atom.


The reaction of obtaining an amine compound of the formula [XVI] from a compound of the formula [XIII] can be conducted, 5 for example, according to a method described in J. Org. Chem., 58, 270 (1993).

Other methods in the above-mentioned scheme can be conducted by ordinary methods.

An alcohol compound of the formula [XII] wherein T is 10 the formula:



can also be produced from a benzyl chloride compound of the formula [XVII] according to the following chemical reaction formula:




15 wherein, R^{10} represents a substituent inactive in Grignard reaction among substituents defined for R^1 .

This reaction is conducted in a solvent, and as the solvent used, ethers such as diethyl ether, tetrahydrofuran, diglyme, 20 triglyme and the like, and mixed solvents of these ethers with aromatic hydrocarbons such as toluene, xylene and the like. If necessary, for example, J. Am. Chem. Soc., 73, 3237 (1951) can be referred.

Among the compounds of the formula [XIII], 25 2-methyl-5-phenylbenzyl chloride of the formula [XIII-1] can

also be produced according to the following scheme.



wherein, L^2 has the same meaning as described above, and R^{11} represents a $\text{C}_1\text{-C}_4$ alkyl group (for example, methyl, ethyl and the like) or vinyl.

The reaction of obtaining a compound of the formula [LIII] from a compound of the formula [LII] can be conducted, for example, according to a method described in Tetrahedron, 50, 13697 (1994).


The reaction of obtaining N,N-dimethyl-(2-methyl-5-phenylbenzyl)amine of the formula [LI] from a compound of the formula [LIII] (Sommelet-Hauser rearrangement) can be conducted, for example, according to a method described in Organic Reactions, 18, 403-464 (1970). As the base used for this reaction, sodium hydride, potassium hydride, sodiumamide, potassiumamide, phenyllithium, butyllithium and the like are listed.

The reaction of obtaining 2-methyl-5-phenylbenzyl chloride of the formula [XIII-1] from N,N-dimethyl-(2-methyl-5-phenyl)benzylamine can be conducted, for example, according to a method described in Tetrahedron

Lett., 24, 3233 (1983).

Production Method B

The compound of the invention of the formula [I] can
5 be produced from a triazolone compound of the formula [XIX]
according to the following chemical reaction formula.



wherein, R¹, L² and T have the same meanings as described above.

The reaction is conducted usually in the presence of
10 a base, and as the base used, for example, inorganic bases such
as sodium hydride, sodium hydroxide, potassium hydroxide,
sodium carbonate, potassium carbonate, and the like are listed.

The reaction temperature of the reaction of obtaining
the compound of the invention of the formula [I] from a triazolone
15 compound of the formula [XIX] is usually in the range from -20
°C to 100°C, and the reaction time is usually in the range from
1 to 100 hours.


Regarding the amounts of reagents used for the reaction,
the ratio of a methylating agent of the formula [IX] is 1 to
20 5 mol and the ratio of the base is 1 to 10 mol, per mol of the
triazolone compound of the formula [XIX].

The reaction is conducted in a solvent if necessary,
and as the solvent used, for example, ethers such as 1,4-dioxane,
tetrahydrofuran, ethylene glycol dimethyl ether, diethylene
25 glycol dimethyl ether, t-butyl methyl ether and the like,
aliphatic hydrocarbons such as hexane, heptane, ligroin,

petroleum ether and the like, aromatic hydrocarbons such as toluene, xylene and the like, organic bases such as pyridine, triethylamine, N-methylaniline, N,N-dimethylaniline, N,N-diethylaniline and the like, nitriles such as acetonitrile, isobutyronitrile and the like, N,N-dimethylformamide, dimethyl sulfoxide, water and the like, and mixtures thereof are listed. These solvents are selected depending on the kind of a base used.

The reaction solution after completion of the reaction is subjected to usual post treatments such as extraction with an organic solvent, concentration and the like, giving isolation of the intended compound. This intended compound can also be purified by re-crystallization, chromatography and the like.

The triazolone compound of the formula [XIX] can be produced, for example, according to the following scheme.



wherein, R^1 , L^2 and T have the same meanings as described above, and M^2 represents sodium or potassium.


An isothiocyanate compound of the formula [XXI] can be produced by reacting a compound of the formula [XIII] with a compound of the formula [XX]. A thiosemicarbazide compound of the formula [XXII] can be produced by reacting the isothiocyanate compound of the formula [XXI] with hydrazine

(anhydrous hydrazine or water-containing hydrazine). A triazolinethion compound of the formula [XXIII] can be produced by reacting the thiosemicarbazide compound of the formula [XXII] with difluoroacetic acid. A sulfonic acid compound of the formula [XXIV] can be produced by oxidizing the triazolinethion compound of the formula [XXIII] using an oxidizing agent (for example, hydrogen peroxide). The triazolone compound of the formula [XIX] can be produced by hydrolyzing the sulfonic acid compound of the formula [XXIV] using an acid (for example, hydrochloric acid). In conducting processes starting from a thiosemicarbazide compound of the formula [XXII] to a triazolone compound of the formula [XIX], for example, JP-A No. 7-196630 can be referred.

The reaction solution after completion of the reaction is subjected to usual post treatments such as extraction with an organic solvent, concentration and the like, giving isolation of the intended compound. This intended compound can also be purified by re-crystallization, chromatography and the like.

20 Production Method C

The compound of the invention of the formula [I] can be produced from a semicarbazide compound of the formula [VII], for example, according to the following chemical reaction formula.



wherein, R^1 and T have the same meanings as described above.

The reaction temperature of the above-mentioned reaction of obtaining the compound of the invention of the formula [I] from a semicarbazide compound of the formula [VII] is usually in the range from -20°C to 100°C, and the reaction time is usually
5 in the range from 1 to 100 hours.

Regarding the amounts of reagents used for the reaction, the ratio of a difluoroacetic anhydride is usually 1 to 5 mol per mol of the semicarbazide compound of the formula [VII].

The reaction is usually conducted in a solvent, and as
10 the solvent used, for example, aromatic hydrocarbons such as toluene, xylene and the like, ethers such as 1,4-dioxane, tetrahydrofuran, diethyl ether and the like, aliphatic hydrocarbons such as hexane, petroleum ether and the like, and mixtures thereof are listed.


15 In this reaction, a difluoroacetyl semicarbazide compound of the formula [XXVI] described later is formed as a by-product in some cases, however, it can be separated off by purifying the intended compound by chromatography and the like. The separated difluoroacetyl semicarbazide compound of
20 the formula [XXVI] can be utilized as a starting material for Production Method F described later.

The reaction solution after completion of the reaction can be, for example after decomposition of excess difluoroacetic anhydride by a base such as an aqueous sodium hydroxide solution
25 and the like, subjected to post treatment operations such as extraction with an organic solvent, concentration and the like, giving isolation of the intended compound. This intended compound can also be purified by re-crystallization, chromatography and the like.

30

Production Method D

The compound of the invention of the formula [I] can be produced from the semicarbazide compound of the formula [VII], for example, according to the following chemical reaction
5 formula.



wherein, R^1 and T have the same meanings as described above, and R^7 represents a C_1 - C_4 alkyl (for example, methyl, ethyl and the like).

10 The reaction temperature of the above-mentioned reaction of obtaining the compound of the invention of the formula [I] from the semicarbazide compound of the formula [VII] is usually in the range from -20°C to 100°C , and the reaction time is usually in the range from 1 to 100 hours.

15 Regarding the amounts of reagents used for the reaction, the ratio of a compound of the formula [XXV] is usually 1 to 100 mol per mol of the semicarbazide compound of the formula [VII].

The reaction is conducted in the presence of a base,
20 and as the base used, for example, metal alkoxides such as sodium methoxide, sodium ethoxide, potassium t-butoxide and the like, alkali metal hydroxides such as sodium hydroxide, potassium hydroxide and the like, alkaline earth metal hydroxides such as calcium hydroxide, magnesium hydroxide and the like, and
25 organic bases such as triethylamine, pyridine and the like are listed, and the ratio of a base is usually 0.1 to 3 mol per

mol of the semicarbazide compound of the formula [VII]


The reaction is conducted in a solvent if necessary, and as the solvent used, for example, alcohols such as methanol, ethanol and the like, ethers such as 1,4-dioxane, tetrahydrofuran, diethyl ether and the like, aliphatic hydrocarbons such as hexane, petroleum ether and the like, aromatic hydrocarbons such as toluene, xylene and the like, amides such as N,N-dimethylformamide and the like, dimethylsulfoxide and the like, and mixtures thereof are listed.

In this reaction, a difluoroacetyl semicarbazide compound of the formula [XXVI] described later is formed as a by-product in some cases, however, it can be separated off by purifying the intended compound by chromatography and the like. The separated difluoroacetyl semicarbazide compound of the formula [XXVI] can be utilized as a starting material for Production Method F described later.

The reaction solution after completion of the reaction can be, after neutralization or concentration if necessary, subjected to post treatment operations such as extraction with an organic solvent, concentration and the like, giving isolation of the intended compound. This intended compound can also be purified by re-crystallization, chromatography and the like.

Production Method E

The compound of the invention of the formula [I] can be produced from a semicarbazide compound of the formula [VII], for example, according to the following chemical reaction formula.



wherein, R¹ and T have the same meanings as described above.

The reaction temperature of the above-mentioned reaction of obtaining the compound of the invention of the formula [I] from the semicarbazide compound of the formula [VII] is usually
 5 in the range from 20°C to 100°C, and the reaction time is usually in the range from 1 to 100 hours.

Regarding the amounts of reagents used for the reaction, the ratio of difluoroacetic acid is usually 1 to 100 mol per
 10 mol of the semicarbazide compound of the formula [VII].

The reaction is conducted in a solvent if necessary, and as the solvent used, for example, aromatic hydrocarbons such as toluene, xylene and the like, aliphatic hydrocarbons such as hexane, petroleum ether and the like, ethers such as
 15 1,4-dioxane, tetrahydrofuran and the like, and mixtures thereof are listed.


In this reaction, the difluoroacetyl semicarbazide compound of the formula [XXVI] described later is formed as a by-product in some cases, however, it can be separated off
 20 by purifying the intended compound by chromatography and the like. The separated difluoroacetyl semicarbazide compound of the formula [XXVI] can be utilized as a starting material for Production Method F described later.

The reaction solution after completion of the reaction
 25 can be, after neutralization or concentration if necessary, subjected to post treatment operations such as extraction with

an organic solvent, concentration and the like, giving isolation of the intended compound. This intended compound can also be purified by re-crystallization, chromatography and the like.

5 Production Method F

The compound of the invention of the formula [I] can be produced from the difluoroacetyl semicarbazide compound of the formula [XXVI], for example, according to the following chemical reaction formula.



wherein, R^1 and T have the same meanings as described above.

The reaction temperature of the reaction of obtaining the compound of the invention of the formula [I] from the difluoroacetyl semicarbazide compound of the formula [XXVI] is usually in the range from 20°C to 100°C, and the reaction time is usually in the range from 10 minutes to 10 hours.


The reaction is conducted in the presence of a base or acid, and as the base used, for example, metal alkoxides such as sodium methoxide, sodium ethoxide, potassium t-butoxide and the like, alkali metal hydroxides such as sodium hydroxide, potassium hydroxide and the like, alkaline earth metal hydroxides such as calcium hydroxide, magnesium hydroxide and the like, and organic bases such as triethylamine, pyridine and the like are listed. The ratio of the base used is usually 0.1 to 10 mol per mol of a difluoroacetyl semicarbazide compound of the formula [XXVI]. As the acid used, for example, organic

acids such as difluoroacetic acid, trifluoroacetic acid and the like are listed, and the ratio of the acid used is usually 1 to 100 mol per mol of the difluoroacetyl semicarbazide compound of the formula [XXVI].

5 The reaction is conducted in a solvent if necessary, and as the solvent used, for example, alcohols such as methanol, ethanol and the like, ethers such as 1,4-dioxane, tetrahydrofuran, diethyl ether and the like, aliphatic hydrocarbons such as hexane, petroleum ether and the like, aromatic hydrocarbons such as toluene, xylene and the like, amides such as N,N-dimethylformamide and the like, sulfoxides such as dimethyl sulfoxide and the like, water and mixtures thereof are listed. These solvents are selected and used depending on the kinds of an acid and base used.

15 The reaction solution after completion of the reaction can be, after neutralization or concentration if necessary, subjected to post treatment operations such as extraction with an organic solvent, concentration and the like, giving isolation of the intended compound. This intended compound can also be purified by re-crystallization, chromatography and the like.

20 The difluoroacetyl semicarbazide compound of the formula [XXVI] can also be obtained as a by-product in Production Methods C, D and E, however, can be produced, for example, according to the following scheme.



wherein, R¹ and T have the same meanings as described above.

The reaction temperature in obtaining the difluoroacetyl semicarbazide compound of the formula [XXVI] from the isocyanate compound of the formula [III] is usually in the range from 0
5 °C to 100°C, and the reaction time is usually in the range from 1 to 10 hours.

The ratio of N'-methyl-difluoroacetohydrazide used for the reaction is usually 1 to 10 mol per mol of the isocyanate compound of the formula [III].


10 The reaction is conducted in a solvent if necessary, and as the solvent used, for example, alcohols such as methanol, ethanol and the like, ethers such as 1,4-dioxane, tetrahydrofuran, diethyl ether and the like, halogenated hydrocarbons such as chloroform, chlorobenzene and the like,
15 aliphatic hydrocarbons such as hexane, petroleum ether and the like, aromatic hydrocarbons such as toluene, xylene and the like, amides such as N,N-dimethylformamide and the like, dimethyl sulfoxide and the like, and mixtures thereof are listed.

The reaction solution after completion of the reaction
20 is subjected, after concentration, to post treatment operations such as extraction with an organic solvent, concentration and the like, giving isolation of the intended compound. This intended compound can also be purified by re-crystallization, chromatography and the like.

25

Production Method G

The compound of the invention of the formula [I] can be produced from a compound of the formula [XIII] and a triazolone compound of the formula [L], for example, according to the
30 following chemical reaction formula.



wherein, R^1 , L^2 and T have the same meanings as described above.

The reaction is conducted in the presence of a base, and as the base used, inorganic bases such as lithium hydroxide, lithium hydride, sodium hydroxide, sodium hydride, potassium hydroxide, potassium hydride, sodium carbonate, potassium carbonate, and the like are listed. The reaction temperature is usually in the range from 0°C to 100°C , and the reaction time is usually in the range from 1 to 100 hours.


Regarding the amounts of reagents used for the reaction, the ratio of the triazolone compound of the formula [L] is usually 1 to 5 mol and the ratio of a base is usually from 1 to 10 mol per mol of the compound of the formula [XIII].

The reaction is conducted in a solvent if necessary, and as the solvent used, for example, alcohols such as methanol, ethanol and the like, ethers such as 1,4-dioxane, ethylene glycol dimethyl ether, diethylene glycol dimethyl ether and the like, aromatic hydrocarbons such as toluene, xylene and the like, organic bases such as pyridine, triethylamine, N-methylaniline, N,N'-dimethylaniline, N,N-diethylaniline and the like, nitriles such as acetonitrile, isobutyronitrile and the like, N,N-dimethylformamide, dimethyl sulfoxide, water and the like, and mixtures thereof are listed. These solvents are selected depending on the kinds of a base used.

The reaction solution after completion of the reaction is subjected to post treatment operations such as extraction

with an organic solvent, concentration and the like, giving isolation of the intended compound. This intended compound can also be purified by re-crystallization, chromatography and the like.

- 5 The triazoline compound of the formula [L] can be produced, for example, according to the following chemical reaction formula.



- 10 The reaction temperature in obtaining the triazolone compound of the formula [L] from 2-methyl semicarbazide is usually in the range from 50°C to 150°C, and the reaction time is usually in the range from 1 to 100 hours.

 The ratio of difluoroacetic acid used for the reaction is usually 1 to 100 mol per mol of 2-methyl semicarbazide.


- 15 The reaction is conducted in a solvent if necessary, and as the solvent used, for example, aromatic hydrocarbons such as toluene, xylene and the like, aliphatic hydrocarbons such as hexane, petroleum ether and the like, ethers such as 1,4-dioxane, tetrahydrofuran and the like, and mixtures thereof
20 are listed.

 In this reaction, 1-difluoroacetyl-2-methyl semicarbazide is formed as a by-product in some cases, however, it can be separated off by purification by chromatography and the like.


- 25 The reaction solution after completion of the reaction can be, after neutralization or concentration if necessary, subjected to post treatment operations such as extraction with

an organic solvent, concentration and the like, giving isolation of the intended compound. This intended compound can also be purified by re-crystallization, chromatography and the like.

The triazolone compound of the formula [L] can be shown
5 as tautomers described below.



2-methyl semicarbazide can be produced, for example, according to the following chemical reaction formula.



10 wherein, M^3 represents sodium or potassium.


This reaction is conducted usually in an acidic aqueous solution (for example, hydrochloric acid).

Trimethylsilyl isocyanate of the above-mentioned formula in which M^3 represents a trimethylsilyl is reacted with
15 methylhydrazine, then, the reaction product is reacted with methanol. Thus, 2-methyl semicarbazide can be produced.

Production Method H

Production can be effected by the following methods (H-1)
20 to (H-6).

(H-1): Method of producing the compound of the invention of the formula [I-2] wherein R^1 in the formula [I] is phenyl, naphthyl or heterocyclic group



wherein, R⁸ represents phenyl, naphthyl or heterocyclic group, and the phenyl, naphthyl or heterocyclic group may optionally be each substituted by one or more substituents selected from the group consisting of halogen atoms, cyano, nitro, C₁-C₁₀ alkyl, C₁-C₁₀ haloalkyl, C₃-C₁₀ cycloalkyl, C₁-C₁₀ alkoxy, C₁-C₁₀ haloalkoxy, C₁-C₁₀ alkylthio, C₁-C₁₀ haloalkylthio, (C₁-C₉ alkyl) carbonyl, (C₁-C₉ alkoxy) carbonyl, (C₁-C₉ alkyl) carbonylamino, phenyl, phenoxy, benzyloxy, C₃-C₂₀ trialkylsilyl, methylenedioxy and difluoromethylenedioxy; X and Z are the same or different and represent a chlorine atom, bromine atom, iodine atom or trifluoromethanesulfonyloxy; and T has the same meaning as described above.

(1) Method of producing a borate compound of the formula [XXVIII]

A borate compound of the formula [XXVIII] can be produced by reacting a compound of the formula [I-1] with bis(pinacolate)diboron in an organic solvent (for example, dimethyl sulfoxide, dimethylformamide) in the presence of a base (for example, potassium acetate, potassium carbonate) and a palladium catalyst (for example, dichloromethane complex of [bis(diphenylphosphino)ferrocene]dichloropalladium (II),

tetrakis(triphenylphosphine)palladium).

The reaction solution after completion of the reaction can be added to water, then, subjected to post treatments such as extraction with an organic solvent, concentration and the like, giving isolation of the intended compound. This intended compound can also be purified by chromatography and the like.

In this reaction, J. Org. Chem., 60, 7508-7510 (1995) can be referred, if necessary.

(2) Method of producing a compound of the invention of the formula [I-2]


The compound of the invention of the formula [I-2] can be produced by reacting the borate compound of the formula [XXVIII] with a compound of the general formula R^8Z in an organic solvent (for example, dimethoxyethane) in the presence of a base (for example, potassium phosphate, potassium carbonate) and a palladium catalyst (for example, dichloromethane complex of [bis(diphenylphosphino)ferrocene]dichloropalladium (II), palladium acetate (II), tetrakis(triphenylphosphine)palladium, mixtures thereof).

A solvent in the reaction solution after completion of the reaction is distilled off, for example, then, the residue is subjected to chromatography, or the reaction solution after completion of the reaction is subjected to post treatments such as extraction with an organic solvent, concentration and the like. Thus the intended compound can be obtained. This intended compound can also be purified by re-crystallization, chromatography and the like.

In this reaction, Tetrahedron Lett., 38, 7645-7648 (1997) can be referred, if necessary.

30

(H-2) Method shown in the following chemical reaction formula
 {method of producing the compound of the invention of the formula
 [I-3] wherein R¹ in the formula [I] is an alkynyl having a triple
 bond at the bonding end)



wherein, A³-C≡C represents a C₂-C₁₀ alkynyl optionally
 substituted having a triple bond at the end. This alkynyl may
 optionally be substituted by one or more substituents selected
 from the group consisting of halogen atoms, cyano, nitro, C₁-C₁₀
 alkoxy, C₁-C₁₀ haloalkoxy, C₁-C₁₀ alkylthio, C₁-C₁₀ haloalkylthio,
 (C₁-C₉ alkyl) carbonyl, (C₁-C₉ alkoxy) carbonyl, (C₁-C₉ alkyl)
 carbonylamino, phenyl, phenoxy, benzyloxy and C₃-C₂₀
 trialkylsilyl; and T and X are the same as described above.


The reaction of obtaining a compound of the invention
 of the formula [I-3] from a compound of the formula [I-1] can
 be conducted, for example, in an aprotic polar solvent (for
 example, acetonitrile, N,N-dimethylformamide) in the presence
 of a base (for example, secondary amine such as diisopropylamine
 and the like, tertiary amine such as triethylamine and the like)
 and a catalyst (for example, combination of triphenylphosphine,
 copper iodide (I) and palladium catalyst such as
 bis(triphenylphosphine)dichloropalladium {PdCl₂(PPh₃)₂} and
 the like). If necessary, WO98/03464, Example 1, Tetrahedron
 Lett., 16, 4467-4470 (1975), Synthesis, 1980, 627-630, or J.
 Org. Chem., 64, 2070-2079 (1999) can be referred.

The reaction solution after completion of the reaction

can be subjected to post treatment operations such as extraction with an organic solvent, concentration and the like, to obtain the intended compound. This intended compound can also be purified by re-crystallization, chromatography and the like.

5

(H-3) Method shown in the following chemical reaction formula {method of producing a compound of the invention of the formula [I-2] wherein R¹ in the formula [I] represents a phenyl, naphthyl or heterocyclic group}



10


wherein, R⁸, T and X are the same as described above.

The reaction of obtaining the compound of the invention of the formula [I-2] from a compound of the formula [I-1] can be conducted, for example, in a mixed solvent of water and diethoxyethane in the presence of a base (for example, inorganic base such as sodium bicarbonate) and a catalyst (for example, tetrakis(triphenyl)phosphinepalladium {Pd(PPh₃)₄}) (If necessary, see description in W096/35669, Beispiel 2), or in a solvent in the presence of tetrabutylammonium bromide, base (for example, inorganic base such as potassium carbonate) and catalyst (for example, palladium acetate). The reaction solution after completion of the reaction can be subjected to post treatment operations such as extraction with an organic solvent, concentration and the like, to obtain the intended compound. This intended compound can also be purified by re-crystallization, chromatography and the like. If necessary,

25

J. Org. Chem., 62, 7170-7173 (1997) can be referred, for example.

(H-4) Method shown in the following scheme {method of producing a compound of the invention of the formula [I-5] wherein R¹ in the formula [I] represents a phenoxy, naphthyloxy or heterocyclic-oxy optionally substituted}



wherein, X and T have the same meanings as described above, and A⁴ represents a phenyl, naphthyl or heterocyclic group, and the phenyl, naphthyl or heterocyclic group may optionally each be substituted by one or more substituents selected from the group consisting of halogen atoms, cyano, nitro, C₁-C₁₀ alkyl, C₁-C₁₀ haloalkyl, C₃-C₁₀ cycloalkyl, C₁-C₁₀ alkoxy, C₁-C₁₀ haloalkoxy, C₁-C₁₀ alkylthio, C₁-C₁₀ haloalkylthio, (C₁-C₉ alkyl) carbonyl, (C₁-C₉ alkoxy) carbonyl, (C₁-C₉ alkyl) carbonylamino, phenyl, phenoxy, benzyloxy, C₃-C₂₀ trialkylsilyl, methylenedioxy and difluoromethylenedioxy.

(1) Method of production from a compound of the formula [I-4]

The reaction of obtaining a compound of the invention of the formula [I-5] from a compound of the formula [I-4] can

be conducted in an organic solvent (for example, methylene chloride) in the presence of a base (for example, organic base such as triethylamine and the like, inorganic base such as sodium bicarbonate and the like) and a catalyst (for example, copper acetate (II)). If necessary, Tetrahedron Lett., 39, 2937-2940 (1998) can be referred.

(2) Method of production from a compound of the formula [I-1]

The reaction of obtaining a compound of the invention of the formula [I-5] from a compound of the formula [I-1] can be conducted in an organic solvent (for example, toluene, ethyl acetate) in the presence of a base (for example, organic base such as triethylamine and the like, inorganic base such as potassium carbonate, cesium carbonate and the like) and a catalyst (for example, copper iodide (I)). The reaction solution after completion of the reaction can be subjected to post treatment operations such as extraction with an organic solvent, concentration and the like, to obtain the intended compound. This intended compound can also be purified by re-crystallization, chromatography and the like.


In this reaction, if necessary, descriptions in J. Amer. Chem. Soc., 119, 10539-10540 (1997), J. Amer. Chem. Soc., 121, 3224-3225 (1999) and J. Amer. Chem. Soc., 121, 4369-4378 (1999) can be referred.

(3) Method of production from a compound of the formula [XXVIII]

The reaction of obtaining a compound of the invention of the formula [I-5] from a compound of the formula [XXVIII] can be conducted in an organic solvent (for example, methylene chloride) in the presence of a base (for example, organic base such as triethylamine and the like, inorganic base such as sodium bicarbonate and the like) and a catalyst (for example, copper

acetate (II)). If necessary, descriptions in Tetrahedron Lett., 39, 2937-2940 (1998) can be referred.


(H-5) Method shown in the following scheme {method of producing compounds of the invention of the formulae [I-7] and [I-8] wherein R^1 in the formula [I] represents a $A^1-ON=CA^2-$ }



wherein, A⁵ represents a group other than a hydrogen atom among represented by A¹, and L², A² and T have the same meanings as described above.

Compounds of the formula [I-7] in which A² is a methyl can be produced from a compound of the formula [I-6] according to a method described in J. Org. Chem., 57, 1481-1486 (1992). Other compounds can also be produced by ordinary methods according to the above-mentioned scheme.

(H-6) Method shown in the following scheme



wherein, A^1 , A^2 , L^2 and T have the same meanings as described above.

Each process in the above-mentioned scheme can be
 5 conducted according to an ordinary method.

When the compound of the invention is used as an active
 ingredient of a fungicidal composition, it may be used itself
 without adding any other components, however, it is usually
 10 mixed with solid carriers, liquid carriers, surfactants and
 other adjuvants, to prepare emulsifiable concentrates, wettable
 powders, water dispersible granules, emulsion formulation,
 flowables, dust formulation, granules and the like, to be used.
 These formulations contain the compound of the invention as
 15 an active ingredient in an amount of usually from 0.1 to 90

wt%.

The solid carrier to be used in the formulation may include, for example, the following materials in fine powder or granule form: mineral materials (e.g., kaolinite clay, attapulgite clay, bentonite clay, montmorillonite clay, acid clay, pyrophyllite, talc, diatomaceous earth, calcite), natural organic materials (e.g., stalks of corn, powder of walnut-shell), synthetic organic materials (e.g., urea), salts (e.g., calcium carbonate, ammonium sulfate) and synthetic inorganic materials (e.g., synthetic hydrated silicon oxide). The liquid carrier may include, for example, aromatic hydrocarbons (e.g., xylene, alkylbenzene, methylnaphthalene), alcohols (e.g., isopropyl alcohol, ethylene glycol, propylene glycol, ethylene glycol mono-ethyl ether), ketones (e.g., acetone, cyclohexanone, isophorone), vegetable oils (e.g., soybean oil, cotton seed oil), petroleum aliphatic hydrocarbons, esters, dimethyl sulfoxide, acetonitrile and water. The surfactant may include, for example, anionic surfactants (e.g., alkylsulfate ester salts, alkyl(aryl)sulfonic acid salts, dialkylsulfosuccinic acid salts, phosphate salts of polyoxyethylene alkyl aryl ether, lignin sulfonic acid salts, naphthalenesulfonic acid formaldehyde condensations), nonionic surfactants (e.g., polyoxyethylene alkyl aryl ethers, polyoxyethylene propylene block copolymer, sorbitan fatty acid esters). The other adjuvants may include, for example, water-soluble polymers (e.g., polyvinyl alcohol, polyvinylpyrrolidone), polysaccharides (e.g., gum arabic, alginic acid and their salts, CMC (carboxy methyl cellulose), xanthan gum), inorganic materials (aluminum magnesium silicate, alumina sol), preservatives, coloring agents, PAP (isopropyl acid phosphate),

BHT.

The method of the application of the compound of the invention is concretely carried out, for example, by a foliar treatment, by a soil treatment, by a seed treatment and the like, also it is carried out by any method which is usually a person in the art uses.

When the compound of the invention is applied as the active ingredient of the fungicidal composition, the application amount of the active ingredient; although it may vary with a variety of plants (crops) to be protected, a variety of plant diseases to be controlled, a extent of disease damage, a formulation type, an application type, application times, weather condition and the like; is usually 1 to 5,000 g per hectare, preferably 5 to 1,000 g per hectare. In the case of emulsifiable concentrates, wettable powders, flowables and the like, they are usually applied after diluted with water, usually at a concentration of 0.0001 to 3% by weight, preferably at a concentration of 0.0005 to 1% by weight as active ingredients of the invention. In the case of dust formulations, granules and the like, they are usually applied as it is without diluting.

The compound of the invention can be used as a agricultural and horticultural antifungal agent in agricultural fields, paddy fields, orchards, tea plantations, grasslands, lawns and the like. Increasing of the antifungal effect may be expected by using the compound of the invention with other agricultural and horticultural antifungal agent(s).

Example of the other agricultural and horticultural antifungal agents include:
azole antifungal compound such as propiconazole, triadimenol, prochloraz, penconazole, tebuconazole, flusilazole,

diniconazole, bromoconazole, epoxyconazole, difenoconazole,
 cyproconazole, metconazole, triflumizole, tetraconazole,
 myclobutanil, fenbuconazole, hexaconazole, fluquinconazole,
 triticonazole, bitertanol, ipconazole, imibenconazole,
 5 imazalil, flutriafol and the like; cyclic amine antifungal
 compound such as fenpropimorph, tridemorph, fenpropidin and
 the like; benzimidazole antifungal compound such as carbendazim,
 benomyl, thiabendazole, thiophanate-methyl and the like;
 ethylen bis(dithiocarbamate) compound such as maneb, zineb,
 10 mancozeb and the like; phthalimide compound such as captan,
 folpet and the like; inorganic copper such as basic copper
 chloride, basic copper sulfate, copper sulfate, copper
 hydroxide and the like; organic copper such as copper-oxinate,
 copper-nonylphenol-sulfonate and the like; procymidone,
 15 cyprodinil, pyrimethanil, diethofencarb, thiuram, fluazinam,
 iprodione, vinclozolin, chlorothalonil, mepanipyrim,
 fenpiclonil, fludioxonil, dichlofluanid, folpet,
 kresoxim-methyl, azoxystrobin, trifloxystrobin,
 picoxystrobin, N-methyl- α -methoxyimino-2-[(2,5-dimethyl-
 20 phenoxy)methyl]phenylacetamide, spiroxamine, quinoxifen,
 fenhexamid, famoxadone, fenamidone (RP-407213), iprovalicarb,
 iminoctadine-triacetate, iminoctadine-albesilate,
 6-iodo-3-propyl-2-propoxy-4(3H)-quinazolinone,
 (Z)-N-(α -cyclopropylmethoxyimino-2,3-difluoro-6-(trifluoro
 25 methyl)benzyl)-2-phenylacetamide, 2-chloro-N-(4'-chloro-
 -(1,1'-biphenyl)-2-yl)-3-pyridinecarboxamide, N-[2-(1,3-
 -dimethylbutyl)-3-thienyl]-2,4-dimethylthiazole-5-
 -carboxamide, N-[2-(1,3-dimethylbutyl)-3-thienyl]-3-
 -trifluoromethyl-1-methylpyrazole-4-carboxamide and the
 30 like.

In the case of using the mixture of the compound of the invention and azole antifungal compound, the ratio of the compound of the invention to azole antifungal compound as the active ingredients is not limited to specific ratio; but when
5 the ratio of azole antifungal compound to the compound of the invention by weight is 0.25 to 4, the excellent synergic effect can be achieved. In the case of using the mixture of the compound of the invention and cyclic amine antifungal compound, the ratio of the compound of the invention to cyclic amine antifungal
10 compound as the active ingredients is not limited to specific ratio; but when the ratio of cyclic amine antifungal compound to the compound of the invention by weight is 1 to 8, the excellent synergic effect can be achieved. In the case of using the mixture of the compound of the invention and quinoxifen, the ratio of
15 the compound of the invention to quinoxifen as the active ingredients is not limited to specific ratio; but when the ratio of quinoxifen to the compound of the invention by weight is 0.25 to 2, the excellent synergic effect can be achieved. In the case of using the mixture of the compound of the invention
20 and spiroxamine, the ratio of the compound of the invention to spiroxamine as the active ingredients is not limited to specific ratio; but when the ratio of spiroxamine to the compound of the invention by weight is 1 to 8, the excellent synergic effect can be achieved. In the case of using the mixture of the compound of the invention and ethylen bis(dithiocarbamate)
25 compound, the ratio of the compound of the invention to ethylen bis(dithiocarbamate) compound as the active ingredients is not limited to specific ratio; but when the ratio of ethylen bis(dithiocarbamate) compound to the compound of the invention
30 by weight is 1 to 20, the excellent synergic effect can be achieved.

In the case of using the mixture of the compound of the invention and phthalimide compound, the ratio of the compound of the invention to phthalimide compound as the active ingredients is not limited to specific ratio; but when the ratio of
5 phthalimide compound to the compound of the invention by weight is 1 to 20, the excellent synergic effect can be achieved. In the case of using the mixture of the compound of the invention and inorganic copper or organic copper, the ratio of the compound of the invention to inorganic copper or organic copper as the
10 active ingredients is not limited to specific ratio; but when the ratio of inorganic copper or organic copper to the compound of the invention by weight is 1 to 20, the excellent synergic effect can be achieved. In the case of using the mixture of the compound of the invention and 6-iodo-3-propyl-2-propoxy-
15 -4(3H)-quinazolinone or (Z)-N-(α -cyclopropylmethoxyimino-2,3-difluoro-6-(trifluoromethyl)benzyl)-2-phenylacetamide, the ratio of the compound of the invention to 6-iodo-3-propyl-2-propoxy-4(3H)-quinazolinone or (Z)-N-(α -cyclopropylmethoxyimino-2,3-difluoro-6-(trifluoromethyl)b
20 enzyl)-2-phenylacetamide as the active ingredients is not limited to specific ratio; when the ratio of 6-iodo-3-propyl-2-propoxy-4(3H)-quinazolinone or (Z)-N-(α -cyclopropylmethoxyimino-2,3-difluoro-6-(trifluoromethyl)b enzyl)-2-phenylacetamide to the compound of the invention by
25 weight is 0.1 to 4, the excellent synergic effect can be achieved. In the case of using the mixture of the compound of the invention and 2-chloro-N-(4'-chloro-(1,1'-biphenyl)-2-yl)-3-piridinecarboxamide, N-[2-(1,3-dimethylbutyl)-3-thienyl]-
-2,4-dimethylthiazole-5-carboxamide or N-[2-(1,3-
30 -dimethylbutyl)-3-thienyl]-3-trifluoromethyl-1-

-methylpyrazole-4-carboxamide, the ratio of the compound of the invention to 2-chloro-N-(4'-chloro-(1,1'-biphenyl)-2-yl)-3-pyridinecarboxamide, N-[2-(1,3-dimethylbutyl)-3-thienyl]-2,4-dimethylthiazole-5-carboxamide or

5 N-[2-(1,3-dimethylbutyl)-3-thienyl]-3-trifluoromethyl-1-methylpyrazole-4-carboxamide as the active ingredients is not limited to specific ratio; but when the ratio of 2-chloro-N-(4'-chloro-(1,1'-biphenyl)-2-yl)-3-pyridinecarboxamide, N-[2-(1,3-dimethylbutyl)-3-thienyl]-2,4-dimethylthiazole-5-carboxamide or N-[2-(1,3-dimethylbutyl)-3-thienyl]-3-trifluoromethyl-1-methylpyrazole-4-carboxamide to the compound of the invention by weight is 0.2 to 5, the excellent synergic effect can be achieved.

6-Iodo-3-propyl-2-propoxy-4(3H)-quinazolinone is a compound described in WO94/26722, (Z)-N-(α -cyclopropylmethoxyimino-2,3-difluoro-6-(trifluoromethyl)benzyl)-2-phenylacetamide is a compound described in WO96/19442, 2-chloro-N-(4'-chloro-(1,1'-biphenyl)-2-yl)-3-pyridinecarboxamide is a compound described in EP545099A, N-[2-(1,3-dimethylbutyl)-3-thienyl]-2,4-dimethylthiazole-5-carboxamide and N-[2-(1,3-dimethylbutyl)-3-thienyl]-3-trifluoromethyl-1-methylpyrazole-4-carboxamide are compounds JP-H09-235282-A.

The compound of the invention can be applied after mixed with other agricultural and horticultural insecticidal agent(s), acaricidal agent(s), nematocidal agent(s), herbicidal agent(s), plant growth controlling agent(s) or fertilizer(s). And also the compound of the invention can be applied at the same time without mixing it beforehand.

30 Examples of the insecticides and acaricides include:

organophosphorus compounds such as fenitrothion [O,O-dimethyl O-(3-methyl-4-nitrophenyl) phosphorothioate], fenthion [O,O-dimethyl O-(3-methyl-4-(methythio)phenyl) phosphorothioate], diazinon [O,O-diethyl O-2-isopropyl-6-methylpyrimidin-4-yl phosphorothioate], chlorpyrifos [O,O-diethyl O-3,5,6-trichloro-2-pyridyl phosphorothioate], acephate [O,S-dimethyl acetylphosphoramidothioate], methidathion [S-2,3-dihydro-5-methoxy-2-oxo-1,3,4-thiadiazol-3-ylmethyl O,O-dimethylphosphorodithioate], disulfoton [O,O-diethyl S-2-ethylthioethylphosphorothioate], DDVP [2,2-dichlorovinyl dimethyl phosphate], Sulprofos [O-ethyl O-4-(methylthio)phenyl S-propyl phosphorodithioate], cyanophos [O-4-cyanophenyl O,O-dimethyl phosphorothioate], dioxabenzofos [2-methoxy-4H-1,3,2-benzodioxaphosphinine-2-sulfide], dimethoate [O,O-dimethyl S-(N-methylcarbamoylmethyl) dithiophosphate], phenthoate [ethyl 2-dimethoxyphosphinothioylthio(phenyl)acetate], malathion [diethyl (dimethoxyphosphinothioylthio)succinate], trichlorfon [dimethyl 2,2,2-trichloro-1-hydroxyethylphosphonate], azinphos-methyl [S-3,4-dihydro-4-oxo-1,2,3-benzotriazin-3-ylmethyl O,O-dimethyl phosphorodithioate], monocrotophos [dimethyl (E)-1-methyl-2-(methylcarbamoyl)-vinylphosphate], ethion [O,O,O',O'-tetraethyl S,S'-methylenebis(phosphorodithioate)], phosphothiazate [N-(O-methyl-S-sec-butyl)phosphorylthiazolidine-2-one] and the like; carbamate compounds such as BPMC (2-sec-butylphenyl methylcarbamate), benfracarb [ethyl N-[2,3-dihydro-2,2-dimethylbenzofuran-7-yloxycarbonyl (methyl)aminothio]-N-isopropyl- β -alaninate], propoxur [2-isopropoxyphenyl N-methylcarbamate], Carbosulfan [2,3-dihydro-2,2-dimethyl-

-7-benzo[b]furanyl N-dibutylaminothio-N-methylcarbamate],
 carbaryl [1-naphthyl-N-methylcarbamate], methomyl
 [S-methyl-N-[(methylcarbamoyl)oxy]thioacetimidate],
 ethiofencarb [2-(ethylthiomethyl)phenylmethylcarbamate],
 5 aldicarb [2-methyl-2-(methylthio)propionaldehyde
 O-methylcarbamoyloxime], oxamyl [N,N-dimethyl-2-methyl-
 carbamoyloxyimino-2-(methylthio)acetamide], fenothiocarb
 [S-4-phenoxybutyl-N,N-dimethylthiocarbamate] and the like;
 pyrethroid compounds such as etofenprox [2-(4-ethoxyphenyl)-
 10 -2-methylpropyl-3-phenoxybenzyl ether], fenvalerate [(RS)-
 α -cyano-3-phenoxybenzyl (RS)-2-(4-chlorophenyl)-3-methyl-
 butyrate], esfenvalerate [(S)- α -cyano-3-phenoxybenzyl
 (S)-2-(4-chlorophenyl)-3-methylbutyrate], fenpropathrin
 [(RS)- α -cyano-3-phenoxybenzyl 2,2,3,3-tetramethyl-
 15 cyclopropanecarboxylate], cypermethrin [(RS)- α -cyano-
 -3-phenoxybenzyl (1RS)-cis,trans-3-(2,2-dichlorovinyl)-
 -2,2-dimethylcyclopropane-carboxylate], permethrin
 [3-phenoxybenzyl (1RS)-cis,trans-3-(2,2-dichlorovinyl)-
 -2,2-dimethylcyclopropanecarboxylate], cyhalothrin [(RS)- α -
 20 -cyano-3-phenoxybenzyl (Z)-(1RS)-cis-3-(2-chloro-3,3,3-
 -trifluoroprop-1-enyl)-2,2-dimethylcyclopropanecarboxylate
], deltamethrin [(S)- α -cyano-3-phenoxybenzyl (1R)-cis-3-
 -(2,2-dibromovinyl)-2,2-dimethylcyclopropanecarboxylate],
 cycloprothrin [(RS)- α -cyano-3-phenoxybenzyl (RS)-2,2-
 25 -dichloro-1-(4-ethoxyphenyl)cyclopropanecarboxylate],
 fluvalinate [α -cyano-3-phenoxybenzyl N-(2-chloro- α , α , α -
 -trifluoro-p-tolyl)-D-valinate], bifenthrin [2-methyl-
 biphenyl-3-ylmethyl (Z)-(1RS)-cis-3-(2-chloro-3,3,3-tri-
 fluoroprop-1-enyl)-2,2-dimethyl-cyclopropanecarboxylate],
 30 acrinathrin [cyano-(3-phenoxyphenyl)methyl [1R-{1 α (S*)-

$3\alpha(Z))$]-2,2-dimethyl-3-[3-oxo-3-(2,2,2-trifluoro-1-(tri-
 fluoromethyl)ethoxy-1-propenyl)cyclopropanecarboxylate],
 2-methyl-2-(4-bromodifluoromethoxyphenyl)propyl 3-phenoxy-
 benzyl ether, tralomeethrin [(S)- α -cyano-3-phenoxybenzyl
 5 (1R-cis)-3-[(1RS)(1,2,2,2-tetrabromoethyl)]-2,2-dimethyl-
 cyclopropanecarboxylate], silafluofen [(4-ethoxyphenyl)-
 {3-(4-fluoro-3-phenoxyphenyl)propyl} dimethylsilane] and the
 like; thiadiazine derivatives such as buprofezin [2-tert-
 -butylimino-3-isopropyl-5-phenyl-1,3,5-triaziazinane-
 10 -4-one]; nitroimidazolidine derivatives;
 nereistoxin derivatives such as cartap [S,S'-(2-dimethyl-
 aminotrimethylene) bis(thiocarbamate)], thiocyclam
 [N,N-dimethyl-1,2,3-trithian-5-ylamine], bensultap
 [S,S'-2-dimethylaminotrimethylene di(benzenethiosulfonate)]
 15 and the like;
 N-cyanoamidine derivatives such as N-cyano-N'-methyl-N'-
 -(6-chloro-3-pyridylmethyl)acetamidine and the like;
 chlorinated hydrocarbons such as endosulfan
 [6,7,8,9,10,10-hexachloro-1,5,5a,6,9,9a-hexahydro-6,9-
 20 -methano-2,4,3-benzodioxathiepine oxide], γ -BHC
 [1,2,3,4,5,6-hexachlorocyclohexane],
 1,1-bis(chlorophenyl)-2,2,2-trichloroethanol and the like;
 benzoylphenylurea compounds such as chlorfluazuron
 [1-(3,5-dichloro-4-(3-chloro-5-trifluoromethylpyridin-2--
 25 yloxy)phenyl)-3-(2,6-difluorobenzoyl)urea], teflubenzuron
 [1-(3,5-dichloro-2,4-difluorophenyl)-3-(2,6-difluoro-
 benzoyl)urea], flufenoxuron [1-(4-(2-chloro-4-trifluoro-
 methylphenoxy)-2-fluorophenyl)-3-(2,6-difluorobenzoyl)urea
] and the like;
 30 formamizine derivatives such as amitraz

[N,N'-[(methylimino)dimethylidene]-di-2,4-xylylidene],
 chlordimeform [N'-(4-chloro-2-methylphenyl)-N,N-dimethyl-
 methanimidamide] and the like; thiourea derivatives such as
 diafenthiuron [N-(2,6-diisopropyl-4-phenoxyphenyl)-N'-
 5 -tert-butylcarbodiimide] and the like; phenylpyrazole
 derivatives; tebfenozide
 [N-tert-butyl-N'-(4-ethylbenzoyl)-3,5-dimethylbenzo-
 hydrazide], bromopropylate [isopropyl 4,4'-dibromobenzylate],
 tetradifon [4-chlorophenyl 2,4,5-trichlorophenylsulfone],
 10 quinomethionate [S,S-6-methylquinoxaline-2,3-diylldithio-
 carbonate], propargate [2-(4-tert-butylphenoxy)cyclohexyl-
 prop-2-yl sulfite], fenbutatin oxide [bis[tris(2-methyl-
 -2-phenylpropyl)tin]oxide], hexythiazox [(4RS,5RS)-5-(4--
 chlorophenyl)-N-chlorohexyl-4-methyl-2-oxo-1,3-thiazolidin
 15 e-3-carboxamide], chlofentezine [3,6-bis(2-chlorophenyl)-
 -1,2,4,5-tetrazine], pyridathioben [2-tert-butyl-5-(4--
 tert-butylbenzylthio)-4-chloropyridazine-3(2H)-one],
 fenpyroximate [tert-butyl (E)-4-[(1,3-dimethyl-5-phenoxy-
 pyrazol-4-yl)methyleneaminooxymethyl]-benzoate],
 20 tebfenpyrad [N-4-tert-butylbenzyl]-4-chloro-3-ethyl-
 -1-methyl-5-pyrazolecarboxamide], polynactin complex
 [tetranactin, dinactin, trinactin], milbemectin, avermectin,
 ivermectin, azadilactin [AZAD], pyrimidifen [5-chloro-N-
 -[2-{4-(2-ethoxyethyl)-2,3-dimethylphenoxy}ethyl]-
 25 -6-ethylpyrimidine-4-amine], pymetrozine [2,3,4,5-
 -tetrahydro-3-oxo-4-[(pyridin-3-yl)methyleneamino]-
 -6-methyl-1,2,4-triazine] and the like.

The compound of the invention can control a variety of
 plant diseases, examples are described below:

30 Blast (*Pyricularia oryzae*), *Helminthosporium* leaf spot

(*Cochliobolus miyabeanus*) and sheath blight (*Rhizoctonia solani*) of rice plant; powdery mildew (*Erysiphe graminis*), scab (*Gibberella zeae*), rust (*Puccinia striiformis*, *P. graminis*, *P. recondita*, *P. hordei*), snow blight (*Typhula* sp.,

5 *Micronectriella nivalis*), loose smut (*Ustilago tritici*, *U. nuda*), bunt (*Tilletia caries*), eyespot (*Pseudocercospora herpotrichoides*), scald (*Rhynchosporium secalis*), leaf blight (*Septoria tritici*) and glume blotch (*Leptosphaeria nodorum*) of barley, wheat, oats and rye; melanose (*Diaporthe citri*),

10 scab (*Elsinoe fawcetti*) and penicillium rot (*Penicillium digitatum*, *P. italicum*) of citrus; blossom blight (*Sclerotinia mali*), canker (*Valsa mali*), powdery mildew (*Podosphaera leucotricha*), *Alternaria* leaf spot (*Alternaria mali*) and scab (*Venturia inaequalis*) of apple; scab (*Venturia nashicola*, *V.*

15 *pirina*), black spot (*Alternaria kikuchiana*) and rust (*Gymnosporangium haraeae*) of pear; brown rot (*Sclerotinia cinerea*), scab (*Cladosporium carpophilum*) and *Phomopsis* rot (*Phomopsis* sp.) of peach; anthracnose (*Elsinoe ampelina*), ripe rot (*Glomerella cingulata*), powdery mildew (*Uncinula necator*),

20 rust (*Phakopsora ampelopsidis*), black rot (*Guignardia bidwellii*) and downy mildew (*Plasmopara viticola*) of grape; anthracnose (*Gloeosporium kaki*) and leaf spot (*Cercospora kaki*, *Mycosphaerella nawae*) of Japanese persimmon; anthracnose (*Colletotrichum lagenarium*), powdery mildew (*Sphaerotheca*

25 *fuliginea*), gummy stem blight (*Mycosphaerella melonis*), stem rot (*Fusarium oxysporum*), downy mildew (*Pseudoperonospora cubensis*), late blight (*Phytophthora* sp.) and damping-off (*Pythium* sp.) of melons and cucumbers; early blight (*Alternaria solani*), leaf mold (*Cladosporium fulvum*) and late blight

30 (*Phytophthora infestans*) of tomato; brown spot (*Phomopsis*

vexans) and powdery mildew (*Erysiphe cichoracearum*) of eggplant; alternaria leaf spot (*Alternaria japonica*) and white spot (*Cercospora brassicae*) of vegetables of Cruciferae; Welsh onion rust (*Puccinia allii*); purple stain (*Cercospora kikuchii*), Sphaceloma scab (*Elsinoe glycines*) and pod and stem blight (*Diaporthe phaseolorum* var. *sojae*) of soybean; kidney bean anthracnose (*Colletotrichum lindemthianum*); early leaf spot (*Cercospora personata*) and leaf spot (*Cercospora arachidicola*) of peanut; pea powdery mildew (*Erysiphe pisi*); early blight (*Alternaria solani*) and late blight (*Phytophthora infestans*) of potato; strawberry powdery mildew (*Sphaerotheca humuli*); net blister blight (*Exobasidium reticulatum*) and white scab (*Elsinoe leucospila*) of tea plant; brown spot (*Alternaria longipes*), powdery mildew (*Erysiphe cichoracearum*), anthracnose (*Colletotrichum tabacum*), downy mildew (*Peronospora tabacina*) and (*Phytophthora nicotianae*) of tobacco; beet leaf spot (*Cercospora beticola*); black spot (*Diplocarpon rosae*) and powdery mildew (*Sphaerotheca pannosa*) of rose; leaf spot (*Septoria chrysanthemi-indici*) and white rust (*Puccinia horiana*) of chrysanthemum; gray mold (*Botrytis cinerea*) and stem rot (*Sclerotinia sclerotiorum*) of various crops and the like.

Examples

The following production examples, formulation examples and test examples and the like describe the invention further in detail below, but do not limit the scope of the invention.

First, production examples of compounds of the invention and production examples of intermediates for the production are shown by Production Examples, Reference Production Examples

and Intermediate Production Examples. The numbers of compounds of the invention are indicated by compound numbers described in Tables 1 to 17 described below.

Production Example 1

5 0.50 g (1.63 mmol) of 5-formyl-2-methyl-4-(2-methyl-5-phenylbenzyl)-2,4-dihydro-3H-1,2,4-triazol-3-one (produced by the same manner as in Reference Production Example 1) was added to 4 ml of chloroform, and to this was added 0.79 g (4.91 mmol) of diethylaminosulfur trifluoride under ice
10 cooling. Then, the mixture was stirred at room temperature overnight. The reaction solution was washed with a 5 wt% sodium bicarbonate solution, then, the organic layer was dried over anhydrous magnesium sulfate, and concentrated to obtain 0.54 g (1.63 mmol) of 5-difluoromethyl-2-methyl-4-(2-methyl-
15 -5-phenylbenzyl)-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 1) having a melting point of 72.7°C.

¹H-NMR(CDCl₃, TMS)

δ (ppm): 7.15-7.55 (8H, m), 6.33 (1H, t), 5.04 (2H, s), 3.54 (3H, s), 2.40 (3H, s)

20 Reference Production Example 1

(1) 117 g (998 mmol) of 2-methylbenzonitrile was dropped to 1000 g of a Na-X type zeolite dried powder (Zeolam type F-9 manufactured by Toyo Corp.; 100 mesh or less) while stirring under air cooling. The resulted mixed was allowed to cool,
25 and after cooled to 50°C or lower, to this was added 213 g (1.33 mol) of bromine, and the mixture was stirred at 80 to 85°C for 1.5 hours, then, cooled to 15°C or lower. This was added into mixture of 100 g of potassium carbonate, 500 ml of water and 2 L of methanol at 10°C or lower. The mixture was stirred for
30 0.5 hours, then, filtrated. The filtrated residue, zeolite

powder was washed with 90% methanol water (2.2 L×2), the filtrate and washed solution were combined, and concentrated until the total liquid quality reached 750 ml. The resulted while crystal was filtrated, and the crystal was washed with water to obtain
5 162 g (826 mmol) of 5-bromo-2-methylbenzonitrile.

¹H-NMR (CDCl₃, TMS)

δ (ppm): 7.68 (1H, d), 7.59 (1H, dd), 7.20 (1H, d), 2.50 (3H, s)

(2) A mixture of 27.4 g (140 mmol) of 5-bromo-2-
10 -methylbenzonitrile, 19.5 g (160 mmol) of dihydroxyphenylborane, 629 mg (2.8 mmol) of palladium acetate (II), 45.1 g (140 mmol) of tetrabutylammonium bromide, 48.4 g (350 mmol) of potassium carbonate and 280 ml of water was stirred at 75°C for 11 hours under nitrogen flow. To the reaction solution was added
15 chloroform, the organic layer was washed with water, dried over anhydrous magnesium sulfate, then, concentrated. The residue was re-crystallized from methanol to give 19.9 g (99.9 mmol) of 2-methyl-5-phenylbenzonitrile.

¹H-NMR (CDCl₃, TMS)

20 δ (ppm): 7.35-7.78 (8H, m), 2.59 (3H, s)

(3) Under nitrogen atmosphere, 126 g (650 mmol) of 2-methyl-5-phenylbenzonitrile was dissolved in 700 ml of anhydrous toluene, and to the resulted solution was dropped 780 ml of a toluene solution of diisobutyl aluminum hydride
25 (1.0 M solution; 780 mmol) at -10 to 0°C, and the mixture was stirred at room temperature overnight. To the reaction solution was dropped 25 ml of ethyl acetate, and this was poured into a mixture of 2.5 L of ice water and 270 ml of concentrated hydrochloric acid. After stirring at 70 to 75°C for 1 hour,
30 the mixture was cooled to room temperature, and separated. The

organic layer was washed with water, saturated sodium bicarbonate solution, then, water, and concentrated to obtain 123 g of 2-methyl-5-phenylbenzaldehyde.

$^1\text{H-NMR}$ (CDCl_3 , TMS)

5 δ (ppm): 10.36 (1H, s), 7.30-8.08 (8H, m), 2.72 (3H, s)

(4) 123 g (625 mmol) of 2-methyl-5-phenylbenzaldehyde was dissolved in a mixture of 1.1 L of tetrahydrofuran and 110 ml of water, and to the resulted solution was dropped, at 15 to 20°C, a solution prepared by dissolving 7.7 g (203 mmol) of sodium boron hydride in 70 ml of a 0.1 wt% sodium hydroxide aqueous solution. After completion of dropping, the mixture was stirred for 4 hours at 15 to 20°C. To this reaction solution was dropped 10 g of glacial acetic acid at 20°C or lower, and this was poured into ice water. The reaction solution was
15 extracted with t-butyl methyl ether, the organic layer was washed with water, saturated sodium bicarbonate solution, then, saturated saline, dried over anhydrous magnesium sulfate, then, concentrated to obtain 121 g of 2-methyl-5-phenylbenzyl alcohol.

20 $^1\text{H-NMR}$ (CDCl_3 , TMS)

δ (ppm): 7.20-7.65 (8H, m), 4.74 (2H, s), 2.38 (3H, s), 1.71 (1H, br)

(5) 14.9 g (75.0 mmol) of 2-methyl-5-phenylbenzyl alcohol was dissolved in 100 ml of anhydrous diethyl ether, and to the
25 resulted solution was dropped 8.12 g (30.0 mmol) of phosphorus tribromide under ice cooling. The mixture was stirred further for 3 hours under ice cooling, then, to the reaction solution was added ice water and t-butyl methyl ether, the mixture was stirred for 30 minutes, then, separated. The organic layer
30 was washed with an about 5 wt% sodium bicarbonate solution and

water sequentially, dried over anhydrous magnesium sulfate, then, concentrated to obtain 19.2 g (73.7 mmol) of 2-methyl-5-phenylbenzyl bromide.

$^1\text{H-NMR}$ (CDCl_3 , TMS)

5 δ (ppm): 7.20–7.65 (8H, m), 4.57 (2H, s), 2.45 (3H, s)

(6) 7.83 g (30 mmol) of 2-methyl-5-phenylbenzyl bromide was dissolved in 90 ml of anhydrous tetrahydrofuran, and to this was added 5.84 g (39 mmol) of silver cyanide, and the mixture was heated under reflux for 2 hours. The reaction solution was filtrated, and the filtrate was concentrated to obtain 7.11 g of crude 2-methyl-5-phenylbenzyl isocyanate.

$^1\text{H-NMR}$ (CDCl_3 , TMS)

δ (ppm): 7.25–7.65 (8H, m), 4.53 (2H, s), 2.37 (3H, s)

(7) 14.9 g of crude 2-methyl-5-phenylbenzyl isocyanate was added to 50 ml of toluene, and to the resulted suspension was added 6.01 g (66.7 mmol) of 2-hydroxyacetohydrazide, and the mixture was stirred at 60°C for 1 hour. After cooling to room temperature, the reaction solution was concentrated, and to the residue was added ethanol to precipitate a crystal, obtaining 13.8 g (44.0 mmol) of 1-hydroxyacetyl-4-(2-methyl-5-phenylbenzyl) semicarbazide.

$^1\text{H-NMR}$ (DMSO-d_6)

δ (ppm): 9.47 (1H, br), 7.10–7.90 (8H, m), 6.83 (1H, br), 5.32 (1H, br), 4.26 (2H, d), 3.91 (2H, d), 2.29 (3H, s)

25 (8) 13.8 g (44.0 mmol) of 1-hydroxyacetyl-4-(2-methyl-5-phenylbenzyl) semicarbazide was added to 100 ml of a 2 wt% potassium hydroxide aqueous solution, and the mixture was stirred at 90°C for 2 hours. After cooling to room temperature, the reaction solution was extracted with ethyl acetate, the organic layer was washed with water, dried over anhydrous

30

magnesium sulfate, then, concentrated. The residue was subjected to silica gel column chromatography (eluant: ethyl acetate), to obtain 12.0 g (40.6 mmol) of 5-hydroxymethyl-4-(2-methyl-5-phenylbenzyl)-2,4-dihydro-3H-1,2,4-triazol-3-one.

¹H-NMR (CDCl₃, TMS)

δ (ppm): 7.15-7.60 (8H, m), 5.17 (1H, t), 5.02 (2H, s), 4.28 (2H, d), 2.40 (3H, s)

(9) 6.29 g (21.3 mmol) of 5-hydroxymethyl-4-(2-methyl-5-phenylbenzyl)-2,4-dihydro-3H-1,2,4-triazol-3-one, 3.53 g (25.6 mmol) of potassium carbonate and 3.63 g (25.6 mmol) of methyl iodide were stirred for 4 hours at 60°C in 38 ml of anhydrous N,N-dimethylformamide. To the reaction solution was added 200 ml of a 5 wt% hydrochloric acid aqueous solution, and this was extracted with ethyl acetate. The organic layer was washed with water, 5 wt% sodium bicarbonate solution, then, saturated saline, dried over anhydrous magnesium sulfate, then, concentrated. The residue was subjected to silica gel column chromatography (eluant: hexane/ethyl acetate = 1/1 (v/v), then, ethyl acetate), to obtain 6.52 g (21.2 mmol) of 5-hydroxymethyl-2-methyl-4-(2-methyl-5-phenylbenzyl)-2,4-dihydro-3H-1,2,4-triazol-3-one.

¹H-NMR (CDCl₃, TMS)

δ (ppm): 7.20-7.65 (8H, m), 5.00 (2H, s), 4.29 (2H, d), 3.48 (3H, s), 2.37 (3H, s)

(10) 6.52 g (21.2 mmol) of 5-hydroxymethyl-2-methyl-4-(2-methyl-5-phenylbenzyl)-2,4-dihydro-3H-1,2,4-triazol-3-one and 14 g of active manganese dioxide were stirred for 6 hours in nitrogen atmosphere under reflux with heating in 63 ml of chloroform. The reaction solution was filtrated through celite,

and the filtrate was concentrated. The residue was subjected to silica gel column chromatography (eluant: hexane:ethyl acetate = 1:1), to obtain 4.32 g (14.1 mmol) of 5-formyl-2-methyl-4-(2-methyl-5-phenylbenzyl)-2,4-dihydro-3H-1,2,4-triazol-3-one.

¹H-NMR(CDCl₃, TMS)

δ (ppm): 9.49 (1H, s), 7.10-7.52 (8H, m), 5.25 (2H, s), 3.63 (3H, s), 2.47 (3H, s)

Production Example 2

5.04 g (16.2 mmol) of 4-(5-bromo-2-methylbenzyl)-5-formyl-2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one (produced in the same manner as in Reference Production Example 2) was dissolved in 40 ml of chloroform, to this was added 7.8 g (48.4 mmol) of diethylaminosulfur trifluoride under ice cooling. Then, the mixture was stirred at room temperature overnight. The reaction solution was washed with a 5 wt% sodium bicarbonate solution, then, the organic layer was dried over anhydrous magnesium sulfate, and concentrated. The residue was subjected to silica gel column chromatography (eluant: hexane/ethyl acetate = 1/2 (v/v)), to obtain 5.04 g (15.2 mmol) of 4-(5-bromo-2-methylbenzyl)-5-difluoromethyl-2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 40) having a melting point of 107.6°C.

¹H-NMR(CDCl₃, TMS)

δ (ppm): 7.32 (1H, dd), 7.06 (1H, brs), 7.05 (1H, d), 6.36 (1H, t), 4.94 (2H, s), 3.55 (3H, s), 2.32 (3H, s)

Reference Production Example 2

(1) 81.7 g (417 mmol) of 5-bromo-2-methylbenzonitrile (produced in the same manner as in (1) of Reference Production

Example 1) was dissolved in 500 ml of anhydrous toluene under nitrogen atmosphere, to the resulted solution was dropped 500 ml (500 mmol) of toluene solution (1.0 M) of diisobutyl aluminum hydride at from -10 to 0°C, and the mixture was stirred at room temperature overnight. To the reaction solution was dropped ethyl acetate, and this was poured into a mixture of 2 L of ice water and 170 ml of concentrated hydrochloric acid. After stirring at 70 to 75°C for 1 hour, then, the mixture was cooled to room temperature, and separated. The organic layer was washed with water, saturated sodium bicarbonate solution, then, water, and concentrated to obtain 80.4 g of 5-bromo-2-methylbenzaldehyde.

¹H-NMR(CDCl₃, TMS)

δ (ppm): 10.2 (1H, s), 7.91 (1H, d), 7.58 (1H, dd), 7.15 (1H, d), 2.62 (3H, s)

(2) 161.2 g (810 mmol) of 5-bromo-2-methylbenzaldehyde was dissolved in 1.5 L of tetrahydrofuran and 150 ml of water, and to the resulted solution was dropped, at 15 to 20°C, a solution prepared by dissolving 10.0 g (264 mmol) of sodium boron hydride in 100 ml of a 0.1 wt% sodium hydroxide aqueous solution. After completion of dropping, the mixture was stirred at 15 to 20°C for 4 hours. To the reaction solution was dropped 12.0 g of glacial acetic acid at 20°C or lower, then, this was poured into ice water. The reaction solution was extracted with t-butyl methyl ether, the organic layer was washed with water, further washed twice with a saturated sodium bicarbonate solution, then, washed with saturated saline, and dried over anhydrous magnesium sulfate, then, concentrated to obtain 168 g of 5-bromo-2-methylbenzyl alcohol.

¹H-NMR(CDCl₃, TMS)

δ (ppm): 7.50 (1H, d), 7.30 (1H, dd), 7.00 (1H, d), 4.62 (2H, d), 2.25 (3H, s), 2.08 (1H, t)

(3) 24.4 g (121 mmol) of 5-bromo-2-methylbenzyl alcohol was dissolved in 200 ml of anhydrous diethyl ether, and to the resulted solution was dropped 16.4 g (60.7 mmol) of phosphorus tribromide under ice cooling. The mixture was stirred further for 3 hours under ice cooling, then, the reaction solution was poured into ice water, then, separated. The organic layer was washed with water, saturated sodium bicarbonate solution and saline sequentially, dried over anhydrous magnesium sulfate, then, concentrated to obtain 29.1 g (110 mmol) of 5-bromo-2-methylbenzyl bromide in the form of brown solid.

$^1\text{H-NMR}$ (CDCl_3 , TMS)

δ (ppm): 7.44 (1H, d), 7.33 (1H, dd), 7.05 (1H, d), 4.43 (2H, s), 2.35 (3H, s)

(4) 23.4 g (88.8 mmol) of 5-bromo-2-methylbenzyl bromide was dissolved in 100 ml of anhydrous tetrahydrofuran, and to this was added 16.0 g (106 mmol) of silver cyanide, and the mixture was heated under reflux for 2 hours. The reaction solution was filtrated, and the filtrate was concentrated to obtain crude 5-bromo-2-methylbenzyl isocyanate. This was mixed with 100 ml of toluene and 8.0 g (88.8 mmol) of 2-hydroxyacetohydrazide, and the mixture was stirred at 60 °C for 2 hours. After cooling to the room temperature, the reaction solution was concentrated, to the residue was added ethanol to precipitate a crystal, obtaining 18.1 g (57.2 mmol) of 4-(5-bromo-2-methylbenzyl)-1-hydroxyacetylsemicarbazide.

$^1\text{H-NMR}$ (DMSO-d_6)

δ (ppm): 9.46 (1H, br), 7.85 (1H, br), 7.02-7.65 (3H, m), 6.85 (1H, br), 5.30 (1H, br), 4.16 (2H, d), 3.90 (2H, d), 2.20 (3H,

s)

(5) 18.0 g (56.9 mmol) of 4-(5-bromo-2-methylbenzyl)-
-1-hydroxyacetyl semicarbazide was added to 170 ml of a 2 wt%
potassium hydroxide aqueous solution, and the mixture was
5 stirred at 90°C for 2 hours. After cooling to the room
temperature, the reaction solution was made acidic by addition
of hydrochloric acid, and the solution was filtrated to obtain
white solid. The filtrate was extracted with ethyl acetate,
the organic layer was washed with water, dried over anhydrous
10 magnesium sulfate, then, concentrated to obtain white solid.
This was combined with the previously obtained white solid and
dried under reduced pressure, to obtain 12.3 g (41.2 mmol) of
4-(5-bromo-2-methylbenzyl)-5-hydroxymethyl-2,4-dihydro-
-3H-1,2,4-triazol-3-one.

15 ¹H-NMR (DMSO-d₆)

δ (ppm): 7.34 (1H, dd), 7.16 (1H, d), 6.95 (1H, d), 5.53 (1H,
br), 4.81 (2H, s), 4.20 (2H, br), 2.27 (3H, s)

(6) 7.16 g (24.0 mmol) of 4-(5-bromo-2-methylbenzyl)-
-5-hydroxymethyl-2,4-dihydro-3H-1,2,4-triazol-3-one, 3.98 g
20 (28.8 mmol) of potassium carbonate and 4.09 g (28.8 mmol) of
methyl iodide were stirred at 60°C for 4 hours in 45 ml of
anhydrous N,N-dimethylformamide. To the reaction solution,
300 ml of a 5 wt% hydrochloric acid aqueous solution was added,
and the mixture was extracted with ethyl acetate. The organic
25 layer was washed with water, 5 wt% sodium bicarbonate solution,
then, saturated saline, and dried over anhydrous magnesium
sulfate, then, concentrated to obtain 7.46 g (23.9 mmol) of
4-(5-bromo-2-methylbenzyl)-5-hydroxymethyl-2-methyl-
-2,4-dihydro-3H-1,2,4-triazol-3-one.

30 ¹H-NMR (CDCl₃, TMS)

δ (ppm): 7.33 (1H, dd), 7.11 (1H, d), 7.05 (1H, d), 4.91 (2H, s), 4.32 (2H, s), 3.49 (3H, s), 2.62 (1H, br), 2.29 (3H, s)
(7) 7.46 g (23.9 mmol) of 4-(5-bromo-2-methylbenzyl)-5-hydroxymethyl-2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one and 15.8 g of active manganese dioxide were stirred for 3 hours in nitrogen atmosphere under reflux with heating in 71 ml of chloroform. The reaction solution was filtrated through celite, and the filtrate was concentrated. The residue was subjected to silica gel column chromatography (eluant: hexane/ethyl acetate = 1/2 (v/v)), to obtain 5.04 g (16.3 mmol) of 4-(5-bromo-2-methylbenzyl)-5-formyl-2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one having a melting point of 107.5°C.

$^1\text{H-NMR}$ (CDCl_3 , TMS)

δ (ppm): 9.49 (1H, s), 7.28 (1H, dd), 7.04 (1H, d), 6.97 (1H, d), 5.15 (2H, s), 3.66 (3H, s), 2.38 (3H, s)

Production Example 3

0.16 g (1.18 mmol) of 4-methylphenylboronic acid, 355 mg (1.07 mmol) of 4-(5-bromo-2-methylbenzyl)-5-difluoromethyl-2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 40; produced by the same method as in Production Example 2), 2 mg (0.008 mmol) of palladium acetate, 0.37 g (2.68 mmol) of potassium carbonate and 0.34 g (1.07 mmol) of tetrabutylammonium bromide were suspended in 5 ml of water, and stirred vigorously under nitrogen flow, then, reacted at a bath temperature of 75°C for 1 hour under nitrogen atmosphere. Water was added the reaction mixture which was then extracted with ethyl acetate, the organic layer was dried over anhydrous magnesium sulfate, then, concentrated. 0.39 g of the residue was subjected to silica gel thin layer chromatography (eluant:

toluene/ethyl acetate = 4/1 (v/v), to obtain 0.33 g of 4-{2-methyl-5-(4-methylphenyl)benzyl}-5-difluoromethyl-2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 4) in the form of oil.

5 $^1\text{H-NMR}(\text{CDCl}_3, \text{TMS})$

δ (ppm): 7.1-7.5 (7H, m), 6.32 (1H, t) 5.03 (2H, s), 3.53 (3H, s), 2.39 (3H, s), 2.38 (3H, s)

Production Example 4

175 mg (1.29 mmol) of 2-fluorophenylboronic acid, 285
10 mg (0.859 mmol) of 4-(5-bromo-2-methylbenzyl)-5-difluoromethyl-2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 40; produced in the same manner as in Production Example 2), 1.06 g of tripotassium phosphate hydrate and 35 mg (0.043 mmol) of [1,1'-
15 -bis(diphenylphosphono)ferrocene]dichloropalladium (II) methylene chloride complex were suspended in 10 ml of 1,2-dimethoxyethane, and stirred for 2 hours in nitrogen atmosphere under reflux with heating. Ethyl acetate was added to the reaction solution and the mixture was filtrated through
20 celite, then, the filtrate was concentrated. 0.43 g of the residue was subjected to silica gel thin layer chromatography (eluant: toluene/ethyl acetate = 4/1 (v/v)), to obtain 0.29 g of 5-difluoromethyl-4-{5-(2-fluorophenyl)-2-methylbenzyl}-2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 18)
25 in the form of oil.

$^1\text{H-NMR}(\text{CDCl}_3, \text{TMS})$

δ (ppm): 7.05-7.45 (7H, m), 6.34 (1H, t), 5.03 (2H, s), 3.54 (3H, s), 2.40 (3H, s)

Production Example 5

30 175 mg (1.29 mmol) of 3-fluorophenylboronic acid, 285

mg (0.859 mmol) of 4-(5-bromo-2-methylbenzyl)-5-difluoromethyl-2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 40; produced in the same manner as in Production Example 2), 1.06 g of tripotassium phosphate hydrate and 35
5 mg (0.043 mmol) of [1,1'-bis(diphenylphosphono)ferrocene]-dichloropalladium (II) methylene chloride complex were suspended in 10 ml of 1,2-dimethoxyethane, and stirred for 2 hours in nitrogen atmosphere under reflux with heating. Ethyl acetate was added to the reaction mixture and the mixture was
10 filtrated through celite, then, the filtrate was concentrated. 0.43 g of the residue was subjected to silica gel thin layer chromatography (eluant: toluene/ethyl acetate = 4/1 (v/v)), to obtain 0.28 g of 5-difluoromethyl-4-{5-(3-fluorophenyl)-2-methylbenzyl}-2-methyl-2,4-dihydro-3H-1,2,4-
15 -triazol-3-one (Compound 19) in the form of oil.

¹H-NMR(CDCl₃, TMS)

δ (ppm): 7.15-7.45 (6H, m), 7.00 (1H, dt), 6.36 (1H, t), 5.04 (2H, s), 3.55 (3H, s), 2.41 (3H, s)

Production Example 6

20 175 mg (1.29 mmol) of 4-fluorophenylboronic acid, 285 mg (0.859 mmol) of 4-(5-bromo-2-methylbenzyl)-5-difluoromethyl-2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 40; produced in the same manner as in Production Example 2), 1.06 g of tripotassium
25 phosphate hydrate and 35 mg (0.043 mmol) of [1,1'-bis(diphenylphosphono)ferrocene]dichloropalladium (II) methylene chloride complex were suspended in 10 ml of 1,2-dimethoxyethane, and stirred for 2 hours in nitrogen atmosphere under reflux with heating. Ethyl acetate was added
30 to the reaction mixture and the mixture was filtrated through

celite, then, the filtrate was concentrated. 0.43 g of the residue was subjected to silica gel thin layer chromatography (eluant: toluene/ethyl acetate = 4/1 (v/v)), to obtain 0.29 g of 5-difluoromethyl-4-{5-(4-fluorophenyl)-2-methylbenzyl}-
5 -2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 20) in the form of oil.

¹H-NMR(CDCl₃, TMS)

δ (ppm): 7.05-7.5 (7H, m), 6.36 (1H, t), 5.03 (2H, s), 3.54 (3H, s), 2.40 (3H, s)

10 Production Example 7

200 mg (0.527 mmol) of 5-difluoromethyl-2-methyl-4-{2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxobororan-2-yl)benzyl}-2,4-dihydro-3H-1,2,4-triazol-3-one (produced in the same manner as in Reference Production Example 3), 124
15 mg (0.683 mmol) of 2-chloro-6-trifluoromethylpyridine, 560 mg (2.64 mmol) of tripotassium phosphate hydrate, 22 mg (0.027 mmol) of {1,1'-bis(diphenylphosphino)ferrocene}-dichloropalladium (II) methylene chloride complex and 3.0 ml of ethylene glycol dimethyl ether were mixed and stirred at
20 80°C for 3 hours. The mixture was cooled, then, ethyl acetate was added to this and the mixture was filtrated through a glass filter precoated with celite. The solvent of the filtrate was distilled off under reduced pressure, and the residue was subjected to silica gel thin layer chromatography {eluant:
25 hexane/ethyl acetate = 1/1 (v/v, addition of 1 vol% methanol)}, to obtain 183 mg of 5-difluoromethyl-2-methyl-4-{2-methyl-5-(6-trifluoromethyl-2-pyridyl)benzyl}-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 329) in the form of pale yellow powder.

30 ¹H-NMR(CDCl₃, TMS)

δ (ppm): 7.8-7.9 (3H, m), 7.72 (1H, s), 7.56 (1H, d), 7.42 (1H, d), 6.36 (1H, t), 5.06 (2H, s), 3.55 (3H, s), 2.44 (3H, s)

Reference Production Example 3

1.00 g (3.01 mmol) of 4-(5-bromo-2-methylbenzyl)-5-difluoromethyl-2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 40; produced in the same manner as in Production Example 2), 0.99 g (3.90 mmol) of bis(pinacolate)diboron, 0.12 g (0.15 mmol) of {1,1'-bis(diphenylphosphino)ferrocene}-dichloropalladium (II) methylene chloride complex, 0.89 g (9.07 mmol) of potassium acetate and 20 ml of dimethyl sulfoxide were mixed, and stirred at 80°C for 8 hours. The mixture was cooled to room temperature, water added to this, and the mixture was extracted with ethyl acetate. The organic layer was washed with water, washed with saturated saline, and dried over anhydrous magnesium sulfate, then, a small amount of hexane was added to this and the mixture was left at 0°C for 3 days. The precipitated crystal was ground, then, filtrated, and washed with hexane, to obtain 1.12 g of 5-difluoromethyl-2-methyl-4-{2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxobororan-2-yl)benzyl}-2,4-dihydro-3H-1,2,4-triazol-3-one in the form of brown powder.

$^1\text{H-NMR}$ (CDCl_3 , TMS)

δ (ppm): 7.66 (1H, d), 7.45 (1H, s), 7.19 (1H, d), 6.21 (1H, t), 4.98 (2H, s), 3.55 (3H, s), 2.34 (3H, s), 1.31 (12H, s)

25 Production Example 8

200 mg (0.527 mmol) of 5-difluoromethyl-2-methyl-4-{2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxobororan-2-yl)benzyl}-2,4-dihydro-3H-1,2,4-triazol-3-one (produced in the same manner as in Reference Production Example 3), 98 mg (0.683 mmol) of 2-chloro-6-methoxypyridine, 560 mg (2.64

mmol) of tripotassium phosphate hydrate, 22 mg (0.027 mmol) of {1,1'-bis(diphenylphosphino)ferrocene}dichloropalladium (II) methylene chloride complex and 3.0 ml of ethylene glycol dimethyl ether were mixed, then stirred at 80°C for 3 hours. 5 The mixture was cooled, then, ethyl acetate was added to this and the mixture was filtrated through a glass filter precoated with celite. The solvent of the filtrate was distilled off under reduced pressure, and the residue was subjected to silica gel thin layer chromatography {eluant: hexane/ethyl acetate = 1/1 (v/v, addition of 1 vol% methanol)}, to obtain 170 mg 10 of 5-difluoromethyl-2-methyl-4-{2-methyl-5-(6-methoxy-2-pyridyl)benzyl}-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 120) in the form of pale yellow liquid.

¹H-NMR (CDCl₃, TMS)

15 δ (ppm): 7.90 (1H, dd), 7.77 (1H, s), 7.58 (1H, t), 7.2-7.3 (2H, m), 6.65 (1H, d), 6.35 (1H, t), 5.05 (2H, s), 3.98 (3H, s), 3.54 (3H, s), 2.42 (3H, s).

Production Example 9

200 mg (0.527 mmol) of 5-difluoromethyl-2-methyl- 20 -4-{2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxobororan-2-yl)benzyl}-2,4-dihydro-3H-1,2,4-triazol-3-one (produced in the same manner as in Reference Production Example 3), 112 mg (0.687 mmol) of 3-bromo-thiophene, 560 mg (2.64 mmol) of tripotassium phosphate hydrate, 22 mg (0.027 mmol) of 25 {1,1'-bis(diphenylphosphino)ferrocene}dichloropalladium (II) methylene chloride complex and 3.0 ml of ethylene glycol dimethyl ether were mixed, and stirred at 80°C for 2 hours. The mixture was cooled, then, ethyl acetate was added to this and the mixture was filtrated through a glass filter precoated with 30 celite. The solvent of the filtrate was distilled off under

reduced pressure, and the residue was subjected to silica gel thin layer chromatography {eluant: hexane/ethyl acetate = 1/1 (v/v, addition of 1 vol% methanol)}, to obtain 146 mg of 5-difluoromethyl-2-methyl-4-{2-methyl-5-(3-thienyl)benzyl}-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 105) in the form of white powder.

¹H-NMR (CDCl₃, TMS)

δ (ppm): 7.42 (1H, dd), 7.2-7.4 (5H, m), 6.33 (1H, t), 5.02 (2H, s), 3.55 (3H, s), 2.38 (3H, s)

10 Production Example 10

200 mg (0.527 mmol) of 5-difluoromethyl-2-methyl-4-{2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxabororan-2-yl)benzyl}-2,4-dihydro-3H-1,2,4-triazol-3-one (produced in the same manner as in Reference Production Example 3), 107 mg (0.793 mmol) of 2-bromo-bis-2-butene, 560 mg (2.64 mmol) of tripotassium phosphate hydrate, 22 mg (0.027 mmol) of {1,1'-bis(diphenylphosphino)ferrocene}dichloropalladium (II) methylene chloride complex and 3.0 ml of ethylene glycol dimethyl ether were mixed, then stirred at 70°C for 2 hours. The mixture was cooled, then, ethyl acetate was added to this and the mixture was filtrated through a glass filter precoated with celite. The solvent of the filtrate was distilled off under reduced pressure, and the residue was subjected to silica gel thin layer chromatography {eluant: hexane/ethyl acetate = 1/1 (v/v, addition of 1 vol% methanol)}, to obtain 123 mg of 5-difluoromethyl-2-methyl-4-{2-methyl-5-((E)-1-methyl-1-propenyl)benzyl}-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 47) in the form of pale yellow powder.

¹H-NMR (CDCl₃, TMS)

δ (ppm): 7.19-7.21 (1H, m), 7.10 (1H, d), 7.00 (1H, s), 6.27

(1H, t), 5.72-7.78 (1H, m), 4.97 (2H, s), 3.54 (3H, s), 2.33 (3H, s), 1.95 (3H, s), 1.76 (3H, d).

Production Example 11

200 mg (0.527 mmol) of 5-difluoromethyl-2-methyl-
5 -4-{2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxobororan-
-2-yl)benzyl}-2,4-dihydro-3H-1,2,4-triazol-3-one (produced
in the same manner as in Reference Production Example 3), 117
mg (0.676 mmol) of 2-bromo-4-methylpyrimidine, 560 mg (2.64
mmol) of tripotassium phosphate hydrate, 22 mg (0.027 mmol)
10 of {1,1'-bis(diphenylphosphino)ferrocene}dichloropalladium
(II) methylene chloride complex and 3.0 ml of ethylene glycol
dimethyl ether were mixed, then stirred at 80°C for 3 hours.
The mixture was cooled, then, ethyl acetate was added to this
and the mixture was filtrated through a glass filter precoated
15 with celite. The solvent of the filtrate was distilled off
under reduced pressure, and the residue was subjected to silica
gel thin layer chromatography (eluant: hexane/ethyl acetate
= 1/1 (v/v, addition of 1 vol% methanol)), to obtain 144 mg
of 5-difluoromethyl-2-methyl-4-{2-methyl-5-(4-methyl-
20 -2-pyrimidinyl)benzyl}-2,4-dihydro-3H-1,2,4-triazol-3-one
(Compound 343) in the form of pale yellow solid.

¹H-NMR(CDCl₃, TMS)

δ (ppm): 8.59 (1H, d), 8.30 (1H, dd), 8.16 (1H, s), 7.30 (1H,
d), 7.01 (1H, d), 6.29 (1H, t), 5.06 (2H, s), 3.56 (3H, s), 2.55
25 (3H, s), 2.41 (3H, s)

Production Example 12

200 mg (0.527 mmol) of 5-difluoromethyl-2-methyl-
-4-{2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxobororan-
-2-yl)benzyl}-2,4-dihydro-3H-1,2,4-triazol-3-one (produced
30 in the same manner as in Reference Production Example 3), 156

mg (0.743 mmol) of 2-iodothiophene, 560 mg (2.64 mmol) of tripotassium phosphate hydrate, 22 mg (0.027 mmol) of {1,1'-bis(diphenylphosphino)ferrocene}dichloropalladium (II) methylene chloride complex and 3.0 ml of ethylene glycol dimethyl ether were mixed, then stirred at 80°C for 2 hours. The mixture was cooled, then, ethyl acetate was added to this and the mixture was filtrated through a glass filter precoated with celite. The solvent of the filtrate was distilled off under reduced pressure, and the residue was subjected to silica gel thin layer chromatography (eluant: hexane/ethyl acetate = 1/1 (v/v, addition of 1 vol% methanol)), to obtain 149 mg of 5-difluoromethyl-2-methyl-4-{2-methyl-5-(2-thienyl)benzyl}-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 104) in the form of white powder.

¹H-NMR(CDCl₃, TMS)

δ (ppm): 7.44 (1H, dd), 7.2-7.3 (4H, m), 7.03-7.06 (1H, m), 6.34 (1H, t), 5.01 (2H, s), 3.55 (3H, s), 2.38 (3H, s)

Production Example 13

200 mg (0.527 mmol) of 5-difluoromethyl-2-methyl-4-{2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxobororan-2-yl)benzyl}-2,4-dihydro-3H-1,2,4-triazol-3-one (produced in the same manner as in Reference Production Example 3), 112 mg (0.683 mmol) of 2-bromothiazole, 560 mg (2.64 mmol) of tripotassium phosphate hydrate, 22 mg (0.027 mmol) of {1,1'-bis(diphenylphosphino)ferrocene}dichloropalladium (II) methylene chloride complex and 3.0 ml of ethylene glycol dimethyl ether were mixed, then stirred at 80°C for 2 hours. The mixture was cooled, then, ethyl acetate was added to this and the mixture was filtrated through a glass filter precoated with celite. The solvent of the filtrate was distilled off

under reduced pressure, and the residue was subjected to silica gel thin layer chromatography (eluant: hexane/ethyl acetate = 1/1 (v/v, addition of 1 vol% methanol)), to obtain 120 mg of 5-difluoromethyl-2-methyl-4-(2-methyl-5-(2-
5 -thiazolyl)benzyl)-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 113) of pale yellow appearance.

¹H-NMR(CDCl₃, TMS)

δ (ppm): 7.78-7.83 (2H, m), 7.63 (1H, s), 7.25-7.29 (2H, m), 6.35 (1H, t), 5.03 (2H, s), 3.56 (3H, s), 2.41 (3H, s)

10 Production Example 14

332 mg (1.00 mmol) of 4-(5-bromo-2-methylbenzyl)-5-difluoromethyl-2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 40), 378 mg (3.00 mmol) of 3-ethyl-3-methoxy-1-pentene, 14 mg (0.020
15 mmol) of PdCl₂(PPh₃)₂, 9.5 mg (0.050 mmol) of copper iodide (I), 15.7 mg (0.060 mmol) of triphenylphosphine and 0.5 ml of triethylamine were added to 4.0 ml of acetonitrile, and the mixture was heated under reflux for 5 hours in nitrogen atmosphere. After cooling to room temperature, t-butyl methyl ether was
20 added to the reaction solution, and the mixture was washed with water and dried over anhydrous magnesium sulfate, then, concentrated under reduced pressure. The residue was subjected to silica gel column chromatography (eluant: hexane/ethyl acetate = 2/1 (v/v)), to obtain 319 mg of
25 5-difluoromethyl-4-(5-(3-ethyl-3-methoxy-1-pentynyl)-2-methylbenzyl)-2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 277).

¹H-NMR(CDCl₃, TMS)

δ (ppm): 7.28 (1H, d), 7.12 (1H, d), 7.02 (1H, s), 6.34 (1H, t),
30 4.95 (2H, s), 3.56 (3H, s), 3.36 (3H, s), 2.36 (3H, s), 1.75

(4H, q), 0.97 (6H, t)

Production Example 15

332 mg (1.00 mmol) of

4-(5-bromo-2-methylbenzyl)-5-difluoromethyl-2-methyl-2,4-
5 -dihydro-3H-1,2,4-triazol-3-one (Compound 40),
332 mg (1.0 mmol) of 3,3-dimethyl-1-butene, 14 mg (0.020 mmol)
of $\text{PdCl}_2(\text{PPh}_3)_2$, 9.5 mg (0.050 mmol) of copper iodide (I), 15.7
mg (0.060 mmol) of triphenylphosphine and 0.5 ml of triethylamine
were added to 4.0 ml of acetonitrile, and the mixture was heated
10 at 40°C for 8 hours in nitrogen atmosphere. After cooling to
room temperature, t-butyl methyl ether was added to the reaction
solution, and the mixture was washed with water and dried over
anhydrous magnesium sulfate, then, concentrated under reduced
pressure. The residue was subjected to silica gel column
15 chromatography (eluant: hexane/ethyl acetate = 2/1 (v/v)), then
to silica gel thin layer chromatography (4 times with
hexane:ethyl acetate = 2:1) to obtain 40 mg (0.13 mmol) of
5-difluoromethyl-4-{5-(3,3-dimethyl-1-butyne)-2-methylben-
zyl}-2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound
20 294).

$^1\text{H-NMR}(\text{CDCl}_3, \text{TMS})$

δ (ppm): 7.22 (1H, d), 7.08 (1H, d), 6.97 (1H, s), 6.31 (1H,
t), 4.93 (2H, s), 3.56 (3H, s), 2.33 (3H, s), 1.29 (9H, s)

Production Example 16

25 664 mg (2.0 mmol) of

4-(5-bromo-2-methylbenzyl)-5-difluoromethyl-2-methyl-2,4-
-dihydro-3H-1,2,4-triazol-3-one (Compound 40),
1.00 g (11.9 mmol) of 2-methyl-3-butyne-2-ol, 70.2 mg (0.10 mmol)
of $\text{PdCl}_2(\text{PPh}_3)_2$, 57.1 mg (0.30 mmol) of copper iodide, 78.7 mg
30 (0.30 mmol) of triphenylphosphine, 1.0 ml of triethylamine and

10 ml of acetonitrile were mixed, and the mixture was heated under reflux for 5 hours in nitrogen atmosphere. After cooling to room temperature, t-butyl methyl ether was added to the reaction solution, and the mixture was filtrated through celite, then, concentrated under reduced pressure. The residue was subjected to silica gel column chromatography (eluant: hexane/ethyl acetate = 2/1 (v/v)) to obtain 400 mg of 5-difluoromethyl-4-{5-(3-hydroxyl-3-methyl-1-butynyl)-2-methylbenzyl}-2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 290).

Production Example 17

3.0 ml of pyridine was added to 400 mg (1.19 mmol) of 5-difluoromethyl-4-{5-(3-hydroxyl-3-methyl-1-butynyl)-2-methylbenzyl}-2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 290), then, 286 mg (2.8 mmol) of acetic anhydride and 12.2 mg (0.10 mmol) of N,N-dimethylaniline were further added. The mixed solution was heated for 7 hours at a bath temperature of 80°C. This mixture was cooled to room temperature, then, poured into a dilute hydrochloric acid-ice mixture. The resulted mixture was extracted twice with ethyl acetate, the organic layer was washed with water, then, with saturated saline, dried over anhydrous magnesium sulfate, then, concentrated under reduced pressure. Hexane was added to the residue and the mixture was stirred, then, filtrated to obtain 295 mg (0.78 mmol) of 4-{5-(3-acetoxy-3-methyl-1-butynyl)-2-methylbenzyl}-5-difluoromethyl-2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 306).

¹H-NMR (CDCl₃, TMS)

δ (ppm): 7.27 (1H, d), 7.12 (1H, d), 7.03 (1H, s), 6.32 (1H, t), 4.94 (2H, s), 3.57 (3H, s), 2.36 (3H, s), 2.07 (3H, s),

1.53)

Production Example 18

498 mg (1.5 mmol) of 4-(5-bromo-2-methylbenzyl)-5-difluoromethyl-2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 40), 378 mg (1.2 mmol) of 3-methyl-3-methoxy-1-butene, 21.1 mg (0.030 mmol) of $\text{PdCl}_2(\text{PPh}_3)_2$, 19.0 mg (0.050 mmol) of copper iodide, 39.3 mg (0.15 mmol) of triphenylphosphine, 607 mg of triethylamine and 6.0 ml of acetonitrile were mixed, and the mixture was heated under reflux for 4.5 hours in nitrogen atmosphere. After cooling to room temperature, t-butyl methyl ether was added to the reaction solution, and the mixture was filtrated through celite, then, concentrated under reduced pressure. The residue was subjected to silica gel column chromatography (eluant: hexane/ethyl acetate = 2/1 (v/v)) to obtain 360 mg of 5-difluoromethyl-2-methyl-4-(2-methyl-5-(3-methyl-methoxy-1-butyryl)-benzyl)-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 326).

$^1\text{H-NMR}$ (CDCl_3 , TMS)

δ (ppm): 7.27 (1H, d), 7.12 (1H, d), 7.01 (1H, s), 6.33 (1H, t), 4.95 (2H, s), 3.40 (3H, s), 2.36 (3H, s), 1.52 (6H, s)

Production Example 19

150 mg (0.396 mmol) of 5-difluoromethyl-2-methyl-4-(2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxabororan-2-yl)benzyl)-2,4-dihydro-3H-1,2,4-triazol-3-one (produced in the same manner as in Reference Production Example 3), 89 mg (0.517 mmol) of 2-bromo-6-methylpyridine, 420 mg (1.98 mmol) of tripotassium phosphate hydrate, 16 mg (0.020 mmol) of {1,1'-bis(diphenylphosphino)ferrocene}dichloropalladium (II) methylene chloride complex and 2.0 ml of ethylene glycol dimethyl ether were mixed, then stirred at 80°C for 3 hours.

The mixture was cooled, then, ethyl acetate was added to this and the mixture was filtrated through a glass filter precoated with celite. The solvent of the filtrate was distilled off under reduced pressure, and the residue was subjected to silica gel thin layer chromatography {eluant: hexane/ethyl acetate = 1/1 (v/v, addition of 1 vol% methanol)}, to obtain 115 mg of 5-difluoromethyl-2-methyl-4-{2-methyl-5-(6-methyl-2-pyridyl)benzyl}-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 117).

¹H-NMR (CDCl₃, TMS)

δ (ppm): 7.84 (1H, d), 7.57-7.34 (2H, m), 7.41 (1H, d), 7.26-7.29 (1H, m), 7.06 (1H, d), 6.30 (1H, t), 5.05 (2H, s), 3.55 (3H, s), 2.59 (3H, s), 2.50 (3H, s)

Production Example 20

150 mg (0.396 mmol) of 5-difluoromethyl-2-methyl-4-{2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxobororan-2-yl)benzyl}-2,4-dihydro-3H-1,2,4-triazol-3-one (produced in the same manner as in Reference Production Example 3), 88 mg (0.595 mmol) of 2,6-dichloropyridine, 420 mg (1.98 mmol) of tripotassium phosphate hydrate, 16 mg (0.020 mmol) of {1,1'-bis(diphenylphosphino)ferrocene}dichloropalladium (II) methylene chloride complex and 2.0 ml of ethylene glycol dimethyl ether were mixed, then stirred at 80°C for 3 hours. The mixture was cooled, then, ethyl acetate was added to this and the mixture was filtrated through a glass filter precoated with celite. The solvent of the filtrate was distilled off under reduced pressure, and the residue was subjected to silica gel thin layer chromatography {eluant: hexane/ethyl acetate = 1/1 (v/v, addition of 1 vol% methanol)}, to obtain 103 mg of 4-{5-(6-chloro-2-pyridyl)-2-methylbenzyl}-5-

-difluoromethyl-2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 118).

¹H-NMR(CDCl₃, TMS)

5 δ (ppm): 7.89 (1H, dd), 7.64-7.70 (2H, m), 7.54 (1H, dd),
7.21-7.31 (2H, m), 6.33 (1H, t), 5.05 (2H, s), 3.56 (3H, t),
2.42 (3H, s)

Production Example 21

150 mg (0.396 mmol) of 5-difluoromethyl-2-methyl-
-4-{2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxobororan-
10 -2-yl)benzyl}-2,4-dihydro-3H-1,2,4-triazol-3-one (produced
in the same manner as in Reference Production Example 3), 94
mg (0.515 mmol) of 2-chloro-4-trifluoromethylpyrimidine, 420
mg (1.98 mmol) of tripotassium phosphate hydrate, 16 mg (0.020
mmol) of {1,1'-bis(diphenylphosphino)ferrocene}-
15 dichloropalladium (II) methylene chloride complex and 2.0 ml
of ethylene glycol dimethyl ether were mixed, then stirred at
80°C for 3 hours. The mixture was cooled, then, ethyl acetate
was added to this and the mixture was filtrated through a glass
filter precoated with celite. The solvent of the filtrate was
20 distilled off under reduced pressure, and the residue was
subjected to silica gel thin layer chromatography (eluant:
hexane/ethyl acetate = 1/1 (v/v, addition of 1 vol% methanol)),
to obtain 132 mg of 5-difluoromethyl-2-methyl-4-{2-methyl-
-5-(4-trifluoromethyl-2-pyrimidinyl)benzyl}-2,4-dihydro-
25 -3H-1,2,4-triazol-3-one (Compound 345).

¹H-NMR(CDCl₃, TMS)

δ (ppm): 8.99 (1H, d), 8.36 (1H, dd), 8.14 (1H, s), 7.46 (1H,
d), 7.34 (1H, d), 6.35 (1H, t), 5.07 (2H, s), 3.57 (3H, s), 2.25
(3H, s)

30 Production Example 22

150 mg (0.396 mmol) of 5-difluoromethyl-2-methyl-
-4-{2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxobororan-
-2-yl)benzyl}-2,4-dihydro-3H-1,2,4-triazol-3-one (produced
in the same manner as in Reference Production Example 3), 116
5 mg (0.516 mmol) of 3-bromo-benzotrifluoride, 420 mg (1.98 mmol)
of tripotassium phosphate hydrate, 16 mg (0.020 mmol) of
{1,1'-bis(diphenylphosphino)ferrocene}dichloropalladium
(II) methylene chloride complex and 2.0 ml of ethylene glycol
dimethyl ether were mixed, then stirred at 80°C for 3 hours.
10 The mixture was cooled, then, ethyl acetate was added to this
and the mixture was filtrated through a glass filter precoated
with celite. The solvent of the filtrate was distilled off
under reduced pressure, and the residue was subjected to silica
gel thin layer chromatography (eluant: hexane/ethyl acetate
15 = 1/1 (v/v, addition of 1 vol% methanol)), to obtain 126 mg
of 5-difluoromethyl-2-methyl-4-{2-methyl-5-(3-
-trifluoromethylphenyl)benzyl}-2,4-dihydro-3H-1,2,4-
-triazol-3-one (Compound 24).

¹H-NMR(CDCl₃, TMS)

20 δ (ppm): 7.73 (1H, s), 7.65-7.67 (1H, m), 7.49-7.59 (2H, m),
7.43 (1H, dd), 7.27-7.30 (1H, m), 7.19 (1H, s), 6.36 (1H, t),
5.05 (2H, s), 3.55 (3H, s), 2.42 (3H, s)

Production Example 23

150 mg (0.396 mmol) of
25 5-difluoromethyl-2-methyl-4-{2-methyl-5-(4,4,5,5-tetra-
methyl-1,3,2-dioxobororan-2-yl)benzyl}-2,4-dihydro-3H-1,2,
4-triazol-3-one (produced in the same manner as in Reference
Production Example 3), 114 mg (0.595 mmol) of
3-bromochlorobenzene, 420 mg (1.98 mmol) of tripotassium
30 phosphate hydrate, 16 mg (0.020 mmol) of {1,1'-bis(diphenyl-

phosphino)ferrocene}dichloropalladium (II) methylene chloride complex and 2.0 ml of ethylene glycol dimethyl ether were mixed, then stirred at 80°C for 3 hours. The mixture was cooled, then, ethyl acetate was added to this and the mixture
5 was filtrated through a glass filter precoated with celite. The solvent of the filtrate was distilled off under reduced pressure, and the residue was subjected to silica gel thin layer chromatography {eluant: hexane/ethyl acetate = 1/1 (v/v, addition of 1 vol% methanol)}, to obtain 118 mg of
10 4-{5-(3-chlorophenyl)-2-methylbenzyl}-5-difluoromethyl-2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 6).
¹H-NMR(CDCl₃, TMS)
δ (ppm): 7.47 (1H, s), 7.25-7.41 (5H, m), 7.15 (1H, s), 6.35 (1H, t), 5.04 (2H, s), 3.55 (3H, s), 2.41 (3H, s)

15 Production Example 24

150 mg (0.396 mmol) of 5-difluoromethyl-2-methyl-4-{2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxobororan-2-yl)benzyl}-2,4-dihydro-3H-1,2,4-triazol-3-one (produced in the same manner as in Reference Production Example 3), 101
20 mg (0.591 mmol) of 3-bromo-toluene, 420 mg (1.98 mmol) of tripotassium phosphate hydrate, 16 mg (0.020 mmol) of {1,1'-bis(diphenylphosphino)ferrocene}dichloropalladium (II) methylene chloride complex and 2.0 ml of ethylene glycol dimethyl ether were mixed, then stirred at 80°C for 3 hours.
25 The mixture was cooled, then, ethyl acetate was added to this and the mixture was filtrated through a glass filter precoated with celite. The solvent of the filtrate was distilled off under reduced pressure, and the residue was subjected to silica gel thin layer chromatography {eluant: hexane/ethyl acetate
30 = 1/1 (v/v, addition of 1 vol% methanol)}, to obtain 106 mg

of 5-difluoromethyl-2-methyl-4-{2-methyl-5-(3-methylphenyl)benzyl}-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 3).

¹H-NMR(CDCl₃, TMS)

5 δ (ppm): 7.41 (1H, dd), 7.23-7.33 (4H, m), 7.13-7.18 (2H, m), 6.32 (1H, t), 5.04 (2H, s), 3.54 (3H, s), 2.40 (2H, s)

Production Example 25

150 mg (0.396 mmol) of 5-difluoromethyl-2-methyl-4-{2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxobororan-2-yl)benzyl}-2,4-dihydro-3H-1,2,4-triazol-3-one (produced in the same manner as in Reference Production Example 3), 114 mg (0.596 mmol) of 4-bromochlorobenzene, 420 mg (1.98 mmol) of tripotassium phosphate hydrate, 16 mg (0.020 mmol) of (1,1'-bis(diphenylphosphino)ferrocene)dichloropalladium (II) methylene chloride complex and 2.0 ml of ethylene glycol dimethyl ether were mixed, then stirred at 80°C for 3 hours. The mixture was cooled, then, ethyl acetate was added to this and the mixture was filtrated through a glass filter precoated with celite. The solvent of the filtrate was distilled off under reduced pressure, and the residue was subjected to silica gel thin layer chromatography (eluant: hexane/ethyl acetate = 1/1 (v/v, addition of 1 vol% methanol)), to obtain 120 mg of 4-{5-(4-chlorophenyl)-2-methylbenzyl}-5-difluoromethyl-2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 7).

25 ¹H-NMR(CDCl₃, TMS)

δ (ppm): 7.35-7.43 (5H, m), 7.24-7.27 (1H, m), 7.15 (1H, s), 6.35 (1H, t), 5.04 (2H, s), 3.54 (3H, s), 2.40 (1H, s)

Production Example 26

300 mg (1.11 mmol) of 2-methyl-4-(2-methyl-5-phenylbenzyl) semicarbazide (produced in the same manner as

in Intermediate Production Example 4) was dissolved in 2 ml of methanol, and to this solution was added, in nitrogen atmosphere, 180 ml (1.45 mmol) of ethyl difluoroacetate and 0.23 ml (1.11 mmol) of a 28% sodium methoxide methanol solution. The mixture was stirred for 9 hours while heating under reflux, cooled to room temperature, then, 20 ml of water was added to the reaction solution, then, 2N hydrochloric acid was added to control pH to 1, and extracted with ethyl acetate. The organic layer was washed with a saturated sodium bicarbonate solution, then with saturated saline, and dried over anhydrous magnesium sulfate, then, concentrated. The residue was subjected to silica gel column chromatography (eluant: hexane/ethyl acetate = 4/1, 2/1, then, 1/2 (v/v)) to obtain 279 mg of 5-difluoromethyl-2-methyl-4-(2-methyl-5-phenylbenzyl)-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 1) and 88 mg of 1-difluoroacetyl-2-methyl-4-(2-methyl-5-phenylbenzyl) semicarbazide.

1-difluoroacetyl-2-methyl-4-(2-methyl-5-phenylbenzyl) semicarbazide:

¹H-NMR (CDCl₃, TMS)

δ (ppm): 8.96 (1H, s), 7.60-7.12 (8H, m), 5.82 (1H, t), 5.45 (1H, t), 4.34 (2H, d), 3.04 (3H, s), 2.26 (3H, s).

Production Example 27

265 mg (0.984 mmol) of 2-methyl-4-(2-methyl-5-phenylbenzyl) semicarbazide (produced in the same manner as in Intermediate Production Example 4) was mixed with 0.5 ml (7.95 mmol) of difluoroacetic acid, and the mixture was stirred at 70°C for 6.5 hours. The reaction solution was cooled to room temperature, then, 5 ml of water was added, and extracted with ethyl acetate. The organic layer was washed with a sodium

bicarbonate solution, then, with saturated saline, and dried over anhydrous magnesium sulfate. The residue was subjected to silica gel column chromatography (eluant: hexane/ethyl acetate = 1/1 (v/v), then ethyl acetate) to obtain 19 mg of
5 5-difluoromethyl-2-methyl-4-(2-methyl-5-phenylbenzyl)-
-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 1) and 297 mg of 1-difluoroacetyl-2-methyl-4-(2-methyl-5-phenylbenzyl) semicarbazide.

Production Example 28

10 146 mg (0.420 mmol) of 1-difluoroacetyl-2-methyl-
-4-(2-methyl-5-phenylbenzyl) semicarbazide (produced in the same manner as in Intermediate Production Example 27) was added to 2 ml of a 2 wt% potassium hydroxide aqueous solution, and the mixture was stirred while heating under reflux for 3 hours.
15 The reaction solution was cooled to room temperature, then, pH was controlled to 3 by addition of 2 N hydrochloric acid, and extracted with ethyl acetate. The organic layer was washed with a sodium bicarbonate solution, then, with saturated saline, and dried over anhydrous magnesium sulfate, then, concentrated
20 to obtain 133 mg of 5-difluoromethyl-2-methyl-4-(2-methyl-
-5-phenylbenzyl)-2,4-dihydro-3H-1,2,4-triazol-3-one
(Compound 1).

Production Example 29

100 mg (0.371 mmol) of
25 2-methyl-4-(2-methyl-5-phenylbenzyl) semicarbazide (produced in the same manner as in Intermediate Production Example 4) and 53 mg (0.557 mmol) of difluoroacetic acid were added to 3.7 ml of toluene, the mixture was stirred at 100°C for 3 hours, then, stirred for 5 hours under reflux with heating. The
30 reaction solution was cooled to room temperature, then, washed

with a saturated sodium bicarbonate solution, then with saturated saline, and dried over anhydrous magnesium sulfate, then, concentrated. The residue was subjected to silica gel thin layer chromatography (eluant: hexane/ethyl acetate = 2/1 (v/v)) to obtain 98 mg of 5-difluoromethyl-2-methyl-4-(2-methyl-5-phenylbenzyl)-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 1) and 14 mg of N-(2-methyl-5-phenylbenzyl)-difluoroacetoamide.

N-(2-methyl-5-phenylbenzyl)-difluoroacetoamide:

¹H-NMR (CDCl₃, TMS)

δ (ppm): 7.19-7.60 (8H, m), 6.57 (1H, br), 5.91 (1H, t), 4.55 (2H, d), 2.35 (3H, s)

Production Example 30

91 mg (0.60 mmol) of 3-methoxyphenylboronic acid, 143 mg (0.43 mmol) of 4-(5-bromo-2-methylbenzyl)-5-difluoromethyl-2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 40; produced in the same manner as in Production Example 2), 0.54 g of tripotassium phosphate hydrate and 18 mg (0.022 mmol) of [1,1'-bis(diphenylphosphono)ferrocene]-dichloropalladium (II) methylene chloride complex were suspended in 8 ml of 1,2-dimethoxyethane, and stirred for 2 hours in nitrogen atmosphere under reflux with heating. Ethyl acetate was added to the reaction solution and the mixture was filtrated through celite, then, the filtrate was concentrated.

The residue (0.19 g) was subjected to silica gel thin layer chromatography (eluant: toluene/ethyl acetate = 4/1 (v/v)), to obtain 0.11 g of 5-difluoromethyl-4-{5-(3-methoxyphenyl)-2-methylbenzyl}-2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 9) in the form of oil.

¹H-NMR (CDCl₃, TMS)

δ (ppm): 6.85-7.45 (7H, m), 6.33 (1H, t), 5.03 (2H, s), 3.84 (3H, s), 3.53 (3H, s), 2.40 (3H, s)

Production Example 31

100 mg (0.671 mmol) of 5-difluoromethyl-2-methyl-
5 -2,4-dihydro-3H-1,2,4-triazol-3-one (produced in the same
manner as in Intermediate Production Example 1) and 175 mg (0.671
mmol) of 2-methyl-5-phenylbenzyl bromide (produced in the same
manner as in Reference Production Example 1 (5)) were dissolved
in 2ml of N,N-dimethylformamide and the solution was cooled
10 to 0°C, and to this was added 40 mg of sodium hydride (1.00
mmol, 60% in oil) slowly. The resulted suspension was allowed
to cool to room temperature, and stirred for 2 hours. The
reaction solution was cooled to 0°C, and water was added slowly,
the mixture was extracted with ethyl acetate. The organic layer
15 was washed with water, saturated sodium bicarbonate solution
and saturated saline sequentially, dried over anhydrous
magnesium sulfate, then, concentrated. The residue was
subjected to silica gel thin layer chromatography (eluant:
hexane/ethyl acetate = 2/1 (v/v)), to obtain 98 mg of
20 5-difluoromethyl-2-methyl-4-(2-methyl-5-phenylbenzyl)-2,4-
-dihydro-3H-1,2,4-triazol-3-one (Compound 1) and 77 mg of
3-difluoromethyl-5-(2-methyl-5-phenylbenzyl)-1-methyl-
-1H-1,2,4-triazole as a by-product.
3-difluoromethyl-5-(2-methyl-5-phenylbenzyl)-1-methyl-1H-
25 -1,2,4-triazole:

¹H-NMR (CDCl₃, TMS)

δ (ppm): 7.28-7.70 (8H, m), 6.54 (1H, t), 5.55 (2H, s), 3.65 (3H, s), 2.45 (3H, s)

Production Example 32

30 5-difluoromethyl-2-methyl-2,4-dihydro-3H-1,2,4-

-triazol-3-one (produced in the same manner as in Intermediate Production Example 1) and 2-methyl-5-phenylbenzyl chloride (produced in the same manner as in Intermediate Production Example 9) were dissolved in N,N-dimethylformamide and the solution was cooled to 0°C, and to this was sodium hydride slowly. The resulting suspension was allowed to cool to room temperature, and stirred. The reaction solution was cooled to 0°C, and water was added slowly, the mixture was extracted with ethyl acetate. The organic layer was washed with water, saturated sodium bicarbonate solution and saturated saline sequentially, dried over anhydrous magnesium sulfate, then, concentrated. The residue was subjected to silica gel thin layer chromatography to obtain 5-difluoromethyl-2-methyl-4-(2-methyl-5-phenylbenzyl)-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 1).

Production Example 33

A mixed suspension of 203 mg (1.36 mmol) of 5-difluoromethyl-2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one (produced in the same manner as in Intermediate Production Example 1), 35 mg (1.46 mmol) of lithium hydroxide and 5 ml of toluene was stirred for 2 hours while heating under reflux. Toluene was distilled off under reduced pressure, 3 ml of anhydrous 1,4-dioxane was added to the residue under nitrogen atmosphere, then, 368 mg (1.33 mmol) of 2-methyl-5-phenylbenzyl methanesulfonate (produced in the same manner as in Intermediate Production Example 11) was added. The reaction solution was stirred for 2 hours while heating under reflux, then, concentrated. Ethyl acetate was added to the residue, the mixture was washed with water, washed with saturated saline, and dried over anhydrous magnesium sulfate, then, concentrated.

The residue was subjected to silica gel thin layer chromatography (eluant: hexane/ethyl acetate = 4/1, then 2/1 (v/v)), to obtain 385 mg of 5-difluoromethyl-2-methyl-4-(2-methyl-5-phenylbenzyl)-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 1) and 31 mg of 3-difluoromethyl-1-methyl-5-(2-methyl-5-phenylbenzyloxy)-1H-1,2,4-triazole as a by-product.

Production Example 34

A mixed suspension of 69 mg (0.461 mmol) of 5-difluoromethyl-2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one (produced in the same manner as in Intermediate Production Example 1), 12 mg (0.508 mmol) of lithium hydroxide, 14 mg (0.092 mmol) of sodium iodide and 3 ml of toluene was stirred for 2 hours while heating under reflux. Toluene was distilled off under reduced pressure, 3 ml of anhydrous 1,4-dioxane was added to the residue under nitrogen atmosphere, then, 100 mg (0.461 mmol) of 2-methyl-5-phenylbenzyl chloride was added. The reaction solution was stirred for 18 hours while heating under reflux, then, concentrated. Ethyl acetate was added to the residue, the mixture was washed with water, then, washed with saturated saline, and dried over anhydrous magnesium sulfate, then, concentrated. The residue was subjected to silica gel thin layer chromatography (eluant: hexane/ethyl acetate = 3/1 (v/v)), to obtain 61 mg of 5-difluoromethyl-2-methyl-4-(2-methyl-5-phenylbenzyl)-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 1) and 11 mg of 3-difluoromethyl-1-methyl-5-(2-methyl-5-phenylbenzyloxy)-1H-1,2,4-triazole as a by-product.

Production Example 35

944 mg (3.35 mmol) of 2-chloro-5-phenylbenzyl bromide (produced in the same manner as in Reference Production Example

5) was dissolved in 10 ml of 1,4-dioxane, and cooled to 0°C. To this solution was added 500 mg (3.35 mmol) of 5-difluoromethyl-2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one (produced in the same manner as in Intermediate Production Example 1) and 40 mg (5.03 mmol) of lithium hydroxide. The mixture was stirred for 30 hours under reflux with heating, then allowed to cool to room temperature. To this mixture was added 6 ml of anhydrous N,N-dimethylformamide, and 180 mg (1.21 mmol) of 5-difluoromethyl-2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one and 200 mg (5.00 mmol, 60%, in oil) of sodium hydride were added, and the mixture was stirred at room temperature for 1 hour, then, a saturated ammonium chloride aqueous solution was added, and extracted with ethyl acetate. The organic layer was washed with a saturated sodium hydrogen carbonate aqueous solution and saturated saline sequentially, dried over anhydrous magnesium sulfate, then, concentrated. The residue was subjected to silica gel column chromatography (eluant: hexane/ethyl acetate = 4/1, then, 2/1 (v/v)) to obtain 824 mg of 5-difluoromethyl-2-methyl-4-(2-chloro-5-phenylbenzyl)-2,4-dihydro-3H-1,2,4-triazol-3-one (Compound 377).

¹H-NMR (CDCl₃, TMS)

δ (ppm): 7.17-7.51 (8H, m), 6.42 (1H, t), 5.18 (2H, s), 3.54 (3H, s)

Reference Production Example 5

(1) A mixture of 5.00 g (24.3 mmol) of 5-bromo-2-chlorotoluene, 3.56 g (29.2 mmol) of dihydroxyphenylborane, 546 mg (2.43 mmol) of palladium acetate (II), 9.41 g (29.2 mmol) of tetrabutylammonium bromide, 8.40 g (60.8 mmol) of potassium carbonate and 50 ml of water was stirred for 5 hours at 80 °C under nitrogen flow. The reaction solution was filtrated

through celite, and the filtrate was extracted with ethyl acetate. The organic layer was washed with saturated saline, dried over anhydrous magnesium sulfate, then, concentrated. The residue was subjected to silica gel column chromatography (eluant:
5 hexane) to obtain 3.92 g of 4-chloro-3-methylbiphenyl.

$^1\text{H-NMR}$ (CDCl_3 , TMS)

δ (ppm): 7.30-7.60 (8H, m), 2.44 (3H, s)

(2) 3.90 g (19.2 mmol) of 4-chloro-3-methylbiphenyl was dissolved in 100 ml of carbon tetrachloride and to this solution
10 was added 632 mg (3.85 mmol) of 2,2'-azobisisobutyronitrile, and the mixture was stirred at room temperature for 15 minutes. To this solution was added 3.43 g (19.2 mmol) of N-bromosuccinimide, and the mixture was stirred for 3.5 hours while heating under reflux. To the reaction solution was
15 further added 632 mg (3.85 mmol) of 2,2'-azobisisobutyronitrile and 300 mg (1.69 mmol) of N-bromosuccinimide, and the mixture was stirred for 2 hours under reflux with heating. The reaction mixture was filtrated, and the filtrate was concentrated. The residue was subjected to silica gel column chromatography
20 (eluant: hexane) to obtain 4.69 g of 2-chloro-5-phenylbenzyl bromide in the form of colorless transparent oil.

$^1\text{H-NMR}$ (CDCl_3 , TMS)

δ (ppm): 7.31-7.69 (8H, m), 4.64 (2H, s)

Next, examples of producing intermediates of the
25 invention are described.

First, the production example of 5-difluoromethyl-2-methyl-2,4-dihydro-3H-1,2,4-triazol-3-one of the formula [L] is shown below.

Intermediate Production Example 1

30 20 ml of toluene was added to 500 mg (5.61 mmol) of 2-methyl

semicarbazide (produced in the same manner as in Reference
Production Example 6), and to this mixture was added 808 mg
(8.41 mmol) of difluoroacetic acid, and the mixture was stirred
for 10 hours under reflux with heating. The reaction solution
5 was cooled to room temperature, then, concentrated, and the
residue was subjected to silica gel thin layer chromatography
(eluant: chloroform/methanol = 9/1 (v/v)), to obtain 305 mg
of 5-difluoromethyl-2-methyl-2,4-dihydro-3H-1,2,4-triazol-
-3-one having a melting point of 141.2°C in the form of colorless
10 solid.

¹H-NMR(CDCl₃, TMS)

δ (ppm): 12.2 (1H,br), 6.46 (1H,t), 3.52 (3H,s)

Next, the production example of difluoroacetyl
semicarbazide of the formula [XXVI] is shown below.

15 Intermediate Production Example 2

1.00 g of crude 2-methyl-5-phenylbenzyl isocyanate was
added to 5 ml of toluene, and to the resulted suspension was
added 556 mg of N'-methyl-difluoroacetohydrazide, and the
mixture was stirred at 60°C. The reaction solution was
20 concentrated, and the residue was subjected to silica gel column
chromatography, to obtain 1-difluoroacetyl-2-methyl-4-(2-
-methyl-5-phenylbenzyl) semicarbazide.

Next, the production example of a semicarbazide compound
of the formula [VII] is shown below.

25 Intermediate Production Example 3

(1) 2.5 g (12.6 mmol) of diphosgene, 0.40 g of activated
carbon and 30 ml of toluene were mixed, and allowed to left
overnight, and to this solution was slowly dropped at room
temperature a solution prepared by dissolving 1.00 g (5.07 mmol)
30 of 2-methyl-5-phenylbenzylamine (produced in the same manner

as in Intermediate Production Example 6) in 5 ml of toluene. The reaction mixture turned to white suspension directly after the dropping. The white suspension was stirred at room temperature for 1 hour, further stirred for 5 hours under reflux
5 with heating, then, concentrated under reduced pressure to obtain 1.13 g of 2-methyl-5-phenylbenzyl isocyanate.

$^1\text{H-NMR}(\text{CDCl}_3, \text{TMS})$

δ (ppm): 7.22-7.65 (8H,m), 4.54 (2H,s), 2.38 (3H,s)

(2) 254 mg (5.52 mmol) of methylhydrazine was dissolved in
10 10 ml of anhydrous toluene, and the mixture was cooled to 0 °C. To this solution was slowly dropped while stirring a solution prepared by dissolving 1.12 g (5.02 mmol) of 2-methyl-5-phenylbenzyl isocyanate in 20 ml of anhydrous toluene. In this operation, the temperature of the reaction
15 solution was controlled in the range from 0 to 5°C. After completion of dropping, the solution was stirred at 0°C for 1 hour, further, stirring was continued for 2 hours at room temperature. The reaction solution was concentrated, and the residue was subjected to silica gel column chromatography
20 (eluant: hexane/ethyl acetate = 1/1, then, 1/2 (v/v), then only ethyl acetate), to obtain 1.28 g (4.75 mmol) of 2-methyl-4-(2-methyl-5-phenylbenzyl) semicarbazide having a melting point of 91.6°C.

$^1\text{H-NMR}(\text{CDCl}_3, \text{TMS})$

25 δ (ppm): 7.18-7.64 (8H,m), 6.68 (1H,br), 4.45 (2H,d), 3.56 (2H,s), 3.16 (3H,s), 2.38 (3H,s)

Intermediate Production Example 4

3.0 g (11.6 mmol) of 2-methyl-5-phenylbenzyl bromide (produced in the same manner as in Reference Production Example
30 1 (5)) was dissolved in 5 ml of anhydrous N,N-dimethylformamide,

and to this solution was added 1.13 g (17.3 mmol) of sodium cyanide, and the mixture was stirred for 3 hours at 100°C, to obtain a 2-methyl-5-phenylbenzyl isocyanate solution. To this solution was added 5 ml of toluene, then, the solution was cooled to 0°C, and to this was slowly dropped 550 mg (11.9 mmol) of methylhydrazine. Further, the mixture was stirred for 3 hours at room temperature, and to this reaction mixture was added water, and the mixture was extracted with chloroform. The organic layer was washed with saturated saline, then, dried over anhydrous magnesium sulfate, and concentrated. The residue was subjected to silica gel column chromatography (eluant: ethyl acetate/methanol = 9/1 (v/v)), to obtain 1.24 g of 2-methyl-4-(2-methyl-5-phenylbenzyl) semicarbazide.

Intermediate Production Example 5

1.0 g (4.61 mmol) of 2-methyl-5-phenylbenzyl chloride (produced in the same manner as in Intermediate Production Example 10) was dissolved in 2 ml of anhydrous N,N-dimethylformamide, and to this solution was added 450 mg (6.92 mmol) of sodium cyanide, and the mixture was stirred at 100°C. Subsequently, 2 ml of toluene was added to the reaction solution and the mixture was cooled to 0°C, to this was added 213 mg (4.61 mmol) of methylhydrazine, the mixture was stirred at room temperature, then, to this was added water, and the mixture was extracted with chloroform. The organic layer was washed with saturated saline, and dried over anhydrous magnesium sulfate, and concentrated to obtain 2-methyl-4-(2-methyl-5-phenylbenzyl) semicarbazide.

Next, the production example of 2-methyl-5-phenylbenzylamine of the formula [XVI-1] is shown below.

Intermediate Production Example 6

1.0 g (5.17 mmol) of 2-methyl-5-phenylbenzonitrile (produced in the same manner as in the above-mentioned Reference Production Example 1 (2)) was dissolved in a mixed solvent of 8 ml of 1,4-dioxane and 8 ml of ethanol, and to this solution was added 1.0 g of a suspension of 2 ml of 28 wt% ammonia water and Raney nickel W2 (manufactured by Aldrich; 50 wt% in water). This mixture was stirred for 8 hours at room temperature under hydrogen atmosphere, then, filtrated through celite, and the celite was washed with ethanol and water. The filtrate and washed solution was combined and concentrated under reduced pressure, and the residue was distilled by using Kugelrohr distillation apparatus (0.5 mmHg, set temperature: 216°C), to obtain 842 mg (4.27 mmol) of 2-methyl-5-phenylbenzylamine.

¹H-NMR (CDCl₃, TMS)

δ (ppm): 7.20-7.66 (8H, m), 3.92 (2H, s), 2.38 (3H, s), 1.47 (2H, brs)

Intermediate Production Example 7

4.46 g (23.1 mmol) of 2-methyl-5-phenylbenzonitrile was dissolved in 26 ml of 1,4-dioxane and to this was added 5 ml of ethanol. To this solution was added 9 ml of 28 wt% ammonia water, and 3 g of a suspension of Raney nickel W2 (manufactured by Aldrich; 50 wt% in water). This mixture was stirred for 8 hours at room temperature under hydrogen atmosphere. To this mixture was further added 2 g of a suspension of Raney nickel W2 (manufactured by Aldrich; 50 wt% in water), 8 ml of 28 wt% ammonia water and 20 ml of ethanol, and the mixture was stirred for 4.5 hours at room temperature under hydrogen atmosphere. The reaction mixture was filtrated through celite, and the filtrate was concentrated. The residue was distilled by using Kugelrohr distillation apparatus (0.5 mmHg, set temperature: 216°C), to

obtain 4.25 g of 2-methyl-5-phenylbenzylamine.

Intermediate Production Example 8

100 mg (0.461 mmol) of 2-methyl-5-phenylbenzyl chloride
(produced in the same manner as in Intermediate Production
5 Example 9) was dissolved in 100 ml of chloroform, and to this
solution was added 65 mg (0.461 mmol) of hexamethyltetramine,
and the mixture was stirred for 15 hours under reflux with heating,
then, 32 mg (0.228 mmol) of hexamethyltetramine was further
added, and stirring was continued for 10 hours under reflux
10 with heating. The resulted crystal was filtrated, and the
crystal was washed with t-butyl methyl ether. The resulted
crystal was dissolved in a mixture of 2 ml of ethanol and 0.5
ml of water, and to this solution was 0.2 ml of concentrated
hydrochloric acid, and the mixture was stirred at room
15 temperature overnight, then, concentrated. To the residue was
added a 2N sodium hydroxide aqueous solution, and the mixture
was extracted with t-butyl methyl ether. The organic layer
was washed with saturated saline, dried over anhydrous magnesium
sulfate, then, the mixture was concentrated to obtain 97 mg
20 of crude 2-methyl-5-phenylbenzylamine.

Next, the production example of 2-methyl-5-phenylbenzyl
chloride of the formula [XIII-1] is shown below.

Intermediate Production Example 9

500 mg (2.52 mmol) of 2-methyl-5-phenylbenzyl alcohol
25 (produced in the same manner as in Reference Production Example
1 (4)) was dissolved in 2 ml of carbon tetrachloride, and to
this solution was added 1.0 g (3.28 mmol) of triphenylphosphine
under nitrogen atmosphere. The reaction mixture was stirred
for 1.5 hours under reflux with heating, cooled to room
30 temperature, then, concentrated. The residue was subjected

to silica gel column chromatography (eluant: hexane/ethyl acetate = 10/1 (v/v)), to obtain 508 mg of 2-methyl-5-phenylbenzyl chloride.

¹H-NMR(CDCl₃, TMS)

5 δ (ppm): 7.19-7.63 (8H, m), 4.67 (2H, s), 2.46 (3H, s)

Intermediate Production Example 10

315 mg (1.40 mmol) of N,N-dimethyl-(2-methyl-5-phenylbenzyl)amine (produced in the same manner as in Intermediate Production Example 12) was dissolved in 3 ml of
10 toluene, and to this solution was added 370 mg (3.92 mmol) of methyl chlorocarbonate at room temperature. Directly after addition, the reaction mixture turned to white suspension. This while suspension was stirred at room temperature for 7 hours, then, washed with 1 N hydrochloric acid, saturated sodium
15 bicarbonate water, and saturated saline, sequentially, and dried over anhydrous magnesium sulfate, then, concentrated. The residue was subjected to silica gel column chromatography (eluant: hexane, then, hexane/ethyl acetate = 10/1 (v/v)), to obtain 282 mg of 2-methyl-5-phenylbenzyl chloride.

20 The production method of a compound of the formula [XIII-2] in which R⁹ is a methyl is described below.

Intermediate Production Example 11

2.00 g (10.1 mmol) of 2-methyl-5-phenylbenzyl alcohol was dissolved in 100 ml of ethyl acetate, and to this solution
25 was added 4.2 ml (30.3 mmol) of triethylamine and the mixture was cooled to 0°C. To the reaction mixture was slowly dropped 1.73 g (15.1 mmol) of methanesulfonyl chloride, and the reaction mixture was controlled to room temperature. After stirring for 1 hours at room temperature, the reaction mixture was made
30 acidic by addition of 1 N hydrochloric acid, and the mixture

was extracted with ethyl acetate. The organic layer was washed with a saturated sodium bicarbonate solution and saturated saline sequentially, dried over anhydrous magnesium sulfate, then, concentrated. The residue was subjected to silica gel
5 column chromatography (eluant: hexane/ethyl acetate = 4/1 (v/v)), to obtain 2.79 g of 2-methyl-5-phenylbenzyl methanesulfonate.

$^1\text{H-NMR}$ (CDCl_3 , TMS)

δ (ppm): 7.25-7.68 (8H, m), 5.34 (2H, s), 2.93 (3H, s), 2.45 (3H, s)

10 Next, the production example of
N-dimethyl-(2-methyl-5-phenylbenzyl)amine is shown below.
Intermediate Production Example 12

(1) A mixture of 3.0 g (14.8 mmol) of 4-phenylbenzyl chloride, 10 ml of toluene and 5.6 ml (26.6 mmol) of a 30 wt%
15 trimethylamine aqueous solution was stirred for 3.7 hours at 60°C. The reaction solution was separated, and the resulted aqueous layer was washed with t-butyl methyl ether. The aqueous layer was filtrated through celite, and the filtrate was concentrated to obtain 4.07 g of (4-methylbenzyl)-
20 trimethylammonium chloride having a melting point of 204.8 °C.

$^1\text{H-NMR}$ ($\text{DMSO}-d_6$, TMS)

δ (ppm): 7.35-7.87 (9H, m), 4.72 (2H, s), 3.11 (9H, s)

(2) 500 mg (1.91 mmol) of (4-phenylbenzyl)-
25 trimethylammonium chloride was dissolved in 20 ml of anhydrous dimethyl sulfoxide, and to the resulted solution was added 153 mg (3.82 mmol, 60% in oil) of sodium hydride at room temperature under nitrogen atmosphere, and the mixture was stirred for 8 hours. To the reaction solution was added 180 ml of saturated
30 saline, and extracted twice with each 70 ml of t-butyl methyl

ether. The organic layer was washed with saturated saline, then, dried over anhydrous magnesium sulfate, and concentrated to obtain 452 mg of N,N-dimethyl-(2-methyl-5-phenylbenzyl)amine.

5 $^1\text{H-NMR}(\text{CDCl}_3, \text{TMS})$

δ (ppm): 7.20-7.66 (8H,m), 3.43 (2H,s), 2.40 (3H,s), 2.27 (6H,s)

Next, the production example of producing a benzyl alcohol compound of the formula [XIII-2] from a benzyl chloride compound of the formula [XVII] is shown below.

10 Reference Production Example 4

264 mg of magnesium and 0.5 ml of anhydrous diethyl ether were charged into a 4-necked round bottomed flask filled with nitrogen, then, small pieces of iodine were placed. After left for a while, the color of iodine disappeared, then, to this
15 was dropped slowly a solution prepared by dissolving 2.0 g (9.87 mmol) of 4-phenylbenzyl chloride in 15 ml of anhydrous diethyl ether. After completion of dropping, the mixture was stirred until most of magnesium was dissolved. Into the resulted brown solution, formaldehyde obtained by thermal decomposition at
20 200°C of 1.48 g (49.4 mmol) of para-formaldehyde sufficiently dried was blown over 1 hour at gas layer part of the reaction vessel together with nitrogen flow. After completion of blowing, stirring was further continued for 2 hours at room temperature, and a saturated ammonium chloride aqueous solution was added.
25 The resulted solution was extracted with t-butyl methyl ether, washed with saturated saline, and concentrated. The residue was subjected to silica gel column chromatography (eluant: hexane/ethyl acetate = 10/1, then 4:1 (v/v)), to obtain 787 mg of 2-methyl-5-phenylbenzyl alcohol.

30 $^1\text{H-NMR}(\text{CDCl}_3, \text{TMS})$

δ (ppm): 7.21-7.63 (8H, m), 4.77 (2H, d), 2.40 (3H, s), 1.56 (1H, dd)

Next, the production example of producing 2-methyl semicarbazide from methylhydrazine is shown below.

5 Reference Production Example 6

6.0 g (130 mmol) of methylhydrazine was dissolved in 25 ml of tetrahydrofuran and cooled to 0°C, to the resulted solution was slowly dropped 11.5 g (100 mmol) of trimethylsilyl isocyanate. After completion of dropping, the mixture was
10 stirred for further 1 hour. Then, 20 ml of methanol was added to the reaction solution, the mixture was stirred for 5 hours at 40°C, and concentrated to obtain 7.77 g of 2-methyl semicarbazide in the form of pale yellow solid.

¹H-NMR(CDCl₃, TMS)

15 δ (ppm): 5.61 (2H, br), 3.80 (2H, brs), 3.15 (3H, s)


Reference Production Example 7

1.3 g (200 mmol) of sodium cyanate, 9.67 g (210 mmol) of methylhydrazine and 100 ml of water were mixed, and the mixture was cooled to 0°C. To the resulted suspension, 18.3 ml (210
20 mmol) of concentrated hydrochloric acid was dropped over 20 minutes, the reaction mixture was controlled to room temperature, then, stirred for 3 hours. Then, the reaction mixture was filtrated, and the filtrate was concentrated to obtain 10.8 g of crude 2-methyl semicarbazide.

25

Next, examples of the compounds of the invention are shown with Compound Number.

The compound of the invention of the formula [I-a] (Table 1 to 15)



5

Table 1

Compound Number	R ¹
1	phenyl
2	2-methylphenyl
3	3-methylphenyl
4	4-methylphenyl
5	2-chlorophenyl
6	3-chlorophenyl
7	4-chlorophenyl
8	2-methoxyphenyl
9	3-methoxyphenyl
10	4-methoxyphenyl
11	2,4-dimethylphenyl
12	3,4-dimethylphenyl
13	2,4-dichlorophenyl
14	3,4-dichlorophenyl
15	4-tert-butylphenyl
16	3-bromophenyl
17	4-bromophenyl
18	2-fluorophenyl
19	3-fluorophenyl
20	4-fluorophenyl
21	4-ethylphenyl
22	3-isopropylphenyl
23	2-trifluoromethylphenyl
24	3-trifluoromethylphenyl
25	4-trifluoromethylphenyl

Table 2

Compound	R ¹
26	3-acetylphenyl
27	4-acetylphenyl
28	3-acetoaminophenyl
29	methyl
30	formyl
31	acetyl
32	butyl
33	tert-butyl
34	hexyl
35	cyclohexyl
36	cyano
37	chloromethyl
38	bromomethyl
39	chloro
40	bromo
41	iodo
42	nitro
43	amino
44	vinyl
45	2-methylpropenyl
46	1,2-dimethylpropenyl
47	(E)-1-methylpropenyl
48	methoxycarbonyl
49	1-pentenyl
50	1-heptenyl

Table 3

Compound	R ¹
51	1-octenyl
52	3-phenylpropenyl
53	benzyl
54	stylyl
55	phenoxy
56	phenylthio
57	2-methylphenoxy
58	3-methylphenoxy
59	4-methylphenoxy
60	2-ethylphenoxy
61	phenoxymethyl
62	phenylthiomethyl
63	benzyloxy
64	benzylthio
65	2-methylphenylthio
66	3-methylphenylthio
67	4-methylphenylthio
68	2-methylbenzyloxy
69	3-methylbenzyloxy
70	4-methylbenzyloxy
71	2-methylbenzylthio
72	3-methylbenzylthio
73	4-methylbenzylthio
74	2-chlorophenoxy
75	3-chlorophenoxy

Table 4

Compound	R ¹
76	4-chlorophenoxy
77	2-chlorophenylthio
78	3-chlorophenylthio
79	4-chlorophenylthio
80	2-methoxyphenylthio
81	3-methoxyphenylthio
82	4-methoxyphenylthio
83	phenoxymethyl
84	2-methylphenoxymethyl
85	3-methylphenoxymethyl
86	4-methylphenoxymethyl
87	3,3-diethylpropynyl
88	2-(trimethylsilyl)ethynyl
89	2-(cyclohexyl)ethynyl
90	3,3-dimethyl3-phenoxypropenyl
91	hydroxymethyl
92	4-nitrophenyl
93	4-cyanophenyl
94	3-cyanophenyl
95	2-cyanophenyl
96	4-ethoxyphenyl
97	4-phenoxyphenyl
98	3-phenoxyphenyl
99	4-trifluoromethoxyphenyl
100	hydroxy

Table 5

Compound	R ¹
101	2-pyridyl
102	3-pyridyl
103	4-pyridyl
104	2-thienyl
105	3-thienyl
106	2-furyl
107	3-furyl
108	2-benzothienyl
109	3-benzothienyl
110	2-pyrimidinyl
111	4-pyrimidinyl
112	1-pyrazolyl
113	2-thiazolyl
114	5-oxazolyl
115	2-quinolyl
116	pyrrol-1-yl
117	6-methyl-2-pyridyl
118	6-chloro-2-pyridyl
119	3-chloro-2-pyridyl
120	6-methoxy-2-pyridyl
121	5-methoxy-2-pyridyl
122	3-methyl-2-pyridyl
123	benzothiazol-2-ylthio
124	3-chloro-5-trifluoromethyl-2-pyridyl
125	2-pyridyloxy

Table 6

Compound	R ⁺
126	3-chloro-5-trifluoromethyl-2-pyridyloxy
127	5-trifluoromethyl-2-pyridyloxy
128	4-pyrimidinyloxy
129	2-pyrimidinyloxy
130	6-chloro-4-pyrimidinyloxy
131	5-chloro-2-pyridyloxy
132	allyloxy
133	propargyloxy
134	cyclopropylmethoxy
135	isopropoxy
136	tert-butoxy
137	N-methylbenzylamino
138	3-chloro-5-trifluoromethyl-2-pyridylamino
139	5-trifluoromethyl-2-pyridylamino
140	4-pyrimidinylthio
141	2-pyrimidinylthio
142	2-pyridylthio
143	5-trifluoromethyl-2-pyridylthio
144	3-chloro-5-trifluoromethyl-2-pyridylthio
145	propylthio
146	tert-butylthio
147	5-methoxy-2-pyridylthio
148	propargylthio
149	cyclopropylmethylthio
150	3,4-dimethylphenylthio

Table 7

Compound	R ¹
151	$\text{CH}_3\text{ON}=\text{C}(\text{CH}_3)$
152	$\text{C}_2\text{H}_5\text{ON}=\text{C}(\text{CH}_3)$
153	$\text{CH}_3\text{CH}_2\text{CH}_2\text{ON}=\text{C}(\text{CH}_3)$
154	$(\text{CH}_3)_2\text{CHON}=\text{C}(\text{CH}_3)$
155	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{ON}=\text{C}(\text{CH}_3)$
156	$(\text{CH}_3)_2\text{CHCH}_2\text{ON}=\text{C}(\text{CH}_3)$
157	$\text{C}_2\text{H}_5(\text{CH}_3)\text{CHON}=\text{C}(\text{CH}_3)$
158	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{ON}=\text{C}(\text{CH}_3)$
159	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{ON}=\text{C}(\text{CH}_3)$
160	$(\text{CH}_3)_3\text{CON}=\text{C}(\text{CH}_3)$
161	1-(cyclopropylmethoxyimino)ethyl
162	1-(cyclopentylmethoxyimino)ethyl
163	$(\text{CH}_3)_3\text{CCH}_2\text{ON}=\text{C}(\text{CH}_3)$
164	$\text{C}_6\text{H}_5\text{CH}_2\text{ON}=\text{C}(\text{CH}_3)$
165	4- $\text{CH}_3\text{C}_6\text{H}_4\text{CH}_2\text{ON}=\text{C}(\text{CH}_3)$
166	3- $\text{CH}_3\text{C}_6\text{H}_4\text{CH}_2\text{ON}=\text{C}(\text{CH}_3)$
167	2- $\text{CH}_3\text{C}_6\text{H}_4\text{CH}_2\text{ON}=\text{C}(\text{CH}_3)$
168	3, 4- $(\text{CH}_3)_2\text{C}_6\text{H}_3\text{CH}_2\text{ON}=\text{C}(\text{CH}_3)$
169	2, 5- $(\text{CH}_3)_2\text{C}_6\text{H}_3\text{CH}_2\text{ON}=\text{C}(\text{CH}_3)$
170	2- $\text{ClC}_6\text{H}_4\text{CH}_2\text{ON}=\text{C}(\text{CH}_3)$
171	3- $\text{ClC}_6\text{H}_4\text{CH}_2\text{ON}=\text{C}(\text{CH}_3)$
172	4- $\text{ClC}_6\text{H}_4\text{CH}_2\text{ON}=\text{C}(\text{CH}_3)$
173	3, 4- $\text{Cl}_2\text{C}_6\text{H}_3\text{CH}_2\text{ON}=\text{C}(\text{CH}_3)$
174	$\text{HC}\equiv\text{CCH}_2\text{ON}=\text{C}(\text{CH}_3)$
175	$\text{H}_2\text{C}=\text{CCH}_2\text{ON}=\text{C}(\text{CH}_3)$

Table 8

Compound	R ¹
176	CH ₃ OCH ₂ CH ₂ ON=C (CH ₃)
177	C ₂ H ₅ OCH ₂ CH ₂ ON=C (CH ₃)
178	C ₆ H ₅ CH ₂ CH ₂ ON=C (CH ₃)
179	2, 4-Cl ₂ C ₆ H ₃ CH ₂ ON=C (CH ₃)
180	C ₆ H ₅ OCH ₂ CH ₂ ON=C (CH ₃)
181	C ₆ H ₅ O (CH ₂) ₃ ON=C (CH ₃)
182	4-CH ₃ C ₆ H ₄ CH ₂ CH ₂ ON=C (CH ₃)
183	2-CH ₃ C ₆ H ₄ CH ₂ CH ₂ ON=C (CH ₃)
184	C ₆ H ₅ O (CH ₂) ₄ ON=C (CH ₃)
185	4-CH ₃ C ₆ H ₄ OCH ₂ CH ₂ ON=C (CH ₃)
186	2-CH ₃ C ₆ H ₄ OCH ₂ CH ₂ ON=C (CH ₃)
187	4-ClC ₆ H ₄ CH ₂ CH ₂ ON=C (CH ₃)
188	2-ClC ₆ H ₄ CH ₂ CH ₂ ON=C (CH ₃)
189	4-CH ₃ C ₆ H ₄ CH (CH ₃) ON=C (CH ₃)
190	2-CH ₃ C ₆ H ₄ CH (CH ₃) ON=C (CH ₃)
191	C ₆ H ₅ CH (CH ₃) ON=C (CH ₃)
192	4- (CH ₃) ₃ CC ₆ H ₄ CH ₂ ON=C (CH ₃)
193	CH ₃ ON=C (CH ₃) CH ₂ ON=C (CH ₃)
194	C ₆ H ₅ ON=C (CH ₃) CH ₂ ON=C (CH ₃)
195	C ₆ H ₅ CH (CN) ON=C (CH ₃)
196	C ₆ H ₅ CH (CH ₃ O) ON=C (CH ₃)
197	C ₆ H ₅ C (=O) O
198	HON=C (CH ₃)
199	C ₆ H ₅ CH ₂ ON=CH
200	5-chloro-2-thienyl

Table 9

Compound	R ¹
201	CH ₃ SC (CH ₃) =N
202	CH ₃ C (=S) NH
203	cyclopropyl-C (=S) NH
204	CH ₃ (CH ₂) ₃ SC (CH ₃) =N
205	CH ₃ SC (C ₂ H ₅) =N
206	CH ₃ SC (CH ₃ CH ₂ CH ₂) =N
207	CH ₃ SC (cyclopropyl) =N
208	C ₆ H ₅ CH ₂ SC (CH ₃) =N
209	C ₆ H ₅ CH ₂ SC (C ₂ H ₅) =N
210	C ₂ H ₅ SC (C ₆ H ₅) =N
211	CH ₃ SC (SCH ₃) =N
212	C ₆ H ₅ SC (SCH ₃) =N
213	C ₆ H ₅ CH ₂ SC (SCH ₃) =N
214	C ₂ H ₅ SC (SC ₂ H ₅) =N
215	C ₆ H ₅ CH ₂ SC (SC ₂ H ₅) =N
216	CH ₃ SC (=S) NH
217	C ₆ H ₅ CH ₂ SC (=S) NH
218	4-CH ₃ C ₆ H ₄ CH ₂ SC (cyclopropyl) =N
219	CH ₃ SC (cyclopentyl) =N
220	CH ₃ SC (cyclohexyl) =N
221	4-CH ₃ OC ₆ H ₄ CH ₂ SC (cyclopropyl) =N
222	4-ClC ₆ H ₄ CH ₂ SC (cyclopropyl) =N
223	C ₆ H ₅ CH ₂ SC (cyclopentyl) =N
224	4-CH ₃ C ₆ H ₄ CH ₂ SC (CH ₃) =N
225	4-ClC ₆ H ₄ CH ₂ SC (CH ₃) =N

Table 10

Compound	R ¹
226	4-CH ₃ OC ₆ H ₄ CH ₂ SC (CH ₃) =N
227	CH ₃ CH ₂ CH ₂ SC (C ₆ H ₅) =N
228	(CH ₃) ₂ CHCH ₂ SC (CH ₃) =N
229	C ₆ H ₅ CH ₂ SC (cyclopropyl) =N
230	4-CH ₃ OC ₆ H ₄ CH ₂ SC (SCH ₃) =N
231	4-ClC ₆ H ₄ CH ₂ SC (SCH ₃) =N
232	4-CH ₃ C ₆ H ₄ CH ₂ SC (SCH ₃) =N
233	4-CF ₃ C ₆ H ₄ CH ₂ SC (SCH ₃) =N
234	4-CF ₃ OC ₆ H ₄ CH ₂ SC (SCH ₃) =N
235	CH ₃ CH ₂ CH ₂ CH ₂ SC (SCH ₃) =N
236	4-ClC ₆ H ₄ CH ₂ SC (=S) NH
237	4-CH ₃ C ₆ H ₄ CH ₂ SC (=S) NH
238	4-CF ₃ C ₆ H ₄ CH ₂ SC (=S) NH
239	2-CH ₃ C ₆ H ₄ CH ₂ SC (SCH ₃) =N
240	4-C ₂ H ₅ C ₆ H ₄ CH ₂ SC (SCH ₃) =N
241	4-ClC ₆ H ₄ CH ₂ SC (S-cyclopropyl) =N
242	4-CH ₃ C ₆ H ₄ CH ₂ SC (S-cyclopropyl) =N
243	4-CF ₃ C ₆ H ₄ CH ₂ SC (S-cyclopropyl) =N
244	4-CF ₃ OC ₆ H ₄ CH ₂ SC (S-cyclopropyl) =N
245	4-CH ₃ OC ₆ H ₄ CH ₂ SC (S-cyclopropyl) =N
246	CH ₃ OCH ₂ CH=C (CH ₃)
247	C ₂ H ₅ OCH ₂ CH=C (CH ₃)
248	C ₆ H ₅ CH=C (CH ₃)
249	C ₆ H ₅ CH ₂ OCH=C (CH ₃)
250	4-CH ₃ C ₆ H ₅ CH ₂ OCH=C (CH ₃)

Table 11

Compound	R ¹
251	$(\text{C}_2\text{H}_5)_3\text{SiC}\equiv\text{C}$
252	$[(\text{CH}_3)_2\text{CH}]_3\text{SiC}\equiv\text{C}$
253	$(\text{CH}_3)_3\text{CSi}(\text{CH}_3)_2\text{C}\equiv\text{C}$
254	$\text{C}_2\text{H}_5\text{Si}(\text{CH}_3)_2\text{C}\equiv\text{C}$
255	$(\text{CH}_3)_2\text{CHSi}(\text{CH}_3)_2\text{C}\equiv\text{C}$
256	$\text{CH}_3\text{Si}(\text{C}_2\text{H}_5)_2\text{C}\equiv\text{C}$
257	$(\text{CH}_3)_2\text{CHSi}(\text{C}_2\text{H}_5)_2\text{C}\equiv\text{C}$
258	$\text{cyclo-C}_3\text{H}_5\text{C}\equiv\text{C}$
259	$\text{cyclopentylC}\equiv\text{C}$
260	$2\text{-ClC}_6\text{H}_4\text{OC}(\text{CH}_3)_2\text{C}\equiv\text{C}$
261	$3\text{-ClC}_6\text{H}_4\text{OC}(\text{CH}_3)_2\text{C}\equiv\text{C}$
262	$4\text{-ClC}_6\text{H}_4\text{OC}(\text{CH}_3)_2\text{C}\equiv\text{C}$
263	$2\text{-CH}_3\text{C}_6\text{H}_4\text{OC}(\text{CH}_3)_2\text{C}\equiv\text{C}$
264	$3\text{-CH}_3\text{C}_6\text{H}_4\text{OC}(\text{CH}_3)_2\text{C}\equiv\text{C}$
265	$4\text{-CH}_3\text{C}_6\text{H}_4\text{OC}(\text{CH}_3)_2\text{C}\equiv\text{C}$
266	$2\text{-CH}_3\text{OC}_6\text{H}_4\text{OC}(\text{CH}_3)_2\text{C}\equiv\text{C}$
267	$3\text{-CH}_3\text{OC}_6\text{H}_4\text{OC}(\text{CH}_3)_2\text{C}\equiv\text{C}$
268	$4\text{-CH}_3\text{OC}_6\text{H}_4\text{OC}(\text{CH}_3)_2\text{C}\equiv\text{C}$
269	$\text{C}_2\text{H}_5\text{C}(\text{CH}_3)_2\text{C}\equiv\text{C}$
270	$\text{CH}_3\text{CH}_2\text{CH}_2\text{C}(\text{CH}_3)_2\text{C}\equiv\text{C}$
271	$(\text{CH}_3)_2\text{CHC}(\text{CH}_3)_2\text{C}\equiv\text{C}$
272	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{C}(\text{CH}_3)_2\text{C}\equiv\text{C}$
273	$(\text{CH}_3)_2\text{CHCH}_2\text{C}(\text{CH}_3)_2\text{C}\equiv\text{C}$
274	$\text{C}_2\text{H}_5\text{CH}(\text{CH}_3)\text{C}(\text{CH}_3)_2\text{C}\equiv\text{C}$
275	$\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{C}(\text{CH}_3)_2\text{C}\equiv\text{C}$

Table 12

Compound	R ¹
276	$\text{CH}_3\text{C}(\text{C}_2\text{H}_5)_2\text{C}\equiv\text{C}$
277	$\text{CH}_3\text{OC}(\text{C}_2\text{H}_5)_2\text{C}\equiv\text{C}$
278	$\text{C}_2\text{H}_5\text{OC}(\text{C}_2\text{H}_5)_2\text{C}\equiv\text{C}$
279	$\text{C}_6\text{H}_5\text{C}(\text{CH}_3)_2\text{C}\equiv\text{C}$
280	$2\text{-CH}_3\text{C}_6\text{H}_4\text{C}(\text{CH}_3)_2\text{C}\equiv\text{C}$
281	$3\text{-CH}_3\text{C}_6\text{H}_4\text{C}(\text{CH}_3)_2\text{C}\equiv\text{C}$
282	$4\text{-CH}_3\text{C}_6\text{H}_4\text{C}(\text{CH}_3)_2\text{C}\equiv\text{C}$
283	$2\text{-ClC}_6\text{H}_4\text{C}(\text{CH}_3)_2\text{C}\equiv\text{C}$
284	$3\text{-ClC}_6\text{H}_4\text{C}(\text{CH}_3)_2\text{C}\equiv\text{C}$
285	$4\text{-ClC}_6\text{H}_4\text{C}(\text{CH}_3)_2\text{C}\equiv\text{C}$
286	$\text{C}_6\text{H}_5\text{C}\equiv\text{C}$
287	$2\text{-CH}_3\text{C}_6\text{H}_4\text{C}\equiv\text{C}$
288	$2\text{-ClC}_6\text{H}_4\text{C}\equiv\text{C}$
289	$2\text{-CH}_3\text{OC}_6\text{H}_4\text{C}\equiv\text{C}$
290	$\text{HOC}(\text{CH}_3)_2\text{C}\equiv\text{C}$
291	$\text{C}_6\text{H}_5\text{CH}_2\text{C}\equiv\text{C}$
292	$\text{C}_6\text{H}_5(\text{CH}_2)_2\text{C}\equiv\text{C}$
293	$\text{C}_6\text{H}_5(\text{CH}_2)_3\text{C}\equiv\text{C}$
294	$(\text{CH}_3)_3\text{CC}\equiv\text{C}$
295	$(\text{CH}_3)_3\text{SiCH}_2\text{C}\equiv\text{C}$
296	$(\text{C}_2\text{H}_5)_3\text{SiCH}_2\text{C}\equiv\text{C}$
297	1-hexynyl
298	$\text{C}_2\text{H}_5\text{OSi}(\text{CH}_3)_2\text{C}\equiv\text{C}$
299	4-methyl-1-pentynyl
300	3-methyl-1-pentynyl

Table 13

Compound	R ¹
301	$\text{C}_6\text{H}_5\text{Si}(\text{CH}_3)_2\text{CH}_2\text{ON}=\text{C}(\text{CH}_3)$
302	$4\text{-CH}_3\text{C}_6\text{H}_4\text{Si}(\text{CH}_3)_2\text{CH}_2\text{ON}=\text{C}(\text{CH}_3)$
303	$2\text{-CH}_3\text{OC}_6\text{H}_4\text{Si}(\text{CH}_3)_2\text{CH}_2\text{ON}=\text{C}(\text{CH}_3)$
304	$2\text{-ClOC}_6\text{H}_4\text{Si}(\text{CH}_3)_2\text{CH}_2\text{ON}=\text{C}(\text{CH}_3)$
305	$\text{C}_6\text{H}_5(\text{CH}_2)_2\text{Si}(\text{CH}_3)_2\text{CH}_2\text{ON}=\text{C}(\text{CH}_3)$
306	$\text{CH}_3\text{COOC}(\text{CH}_3)_2\text{C}\equiv\text{C}$
307	1-cyclohexenyl
308	3-methyl-1-cyclohexenyl
309	$\text{CH}_3\text{O}(\text{CH}_2)_5\text{C}\equiv\text{C}$
310	$\text{C}_2\text{H}_5\text{O}(\text{CH}_2)_5\text{C}\equiv\text{C}$
311	$\text{CH}_3\text{O}(\text{CH}_2)_4\text{C}\equiv\text{C}$
312	$\text{C}_2\text{H}_5\text{O}(\text{CH}_2)_4\text{C}\equiv\text{C}$
313	$\text{CH}_3\text{O}(\text{CH}_2)_3\text{C}\equiv\text{C}$
314	$\text{C}_2\text{H}_5\text{O}(\text{CH}_2)_3\text{C}\equiv\text{C}$
315	cyclo- $\text{C}_3\text{H}_5\text{O}(\text{CH}_2)_3\text{C}\equiv\text{C}$
316	cyclo- $\text{C}_3\text{H}_5\text{C}(\text{CH}_3)(\text{CH}_3\text{O})\text{C}\equiv\text{C}$
317	cyclo- $\text{C}_3\text{H}_5\text{C}(\text{CH}_3)(\text{C}_2\text{H}_5\text{O})\text{C}\equiv\text{C}$
318	cyclo- $\text{C}_3\text{H}_5\text{OC}(\text{CH}_3)(\text{cyclo-}\text{C}_3\text{H}_5)\text{C}\equiv\text{C}$
319	$\text{C}_2\text{H}_5\text{C}(\text{CH}_3)(\text{CH}_3\text{O})\text{C}\equiv\text{C}$
320	$\text{C}_2\text{H}_5\text{C}(\text{CH}_3)(\text{C}_2\text{H}_5\text{O})\text{C}\equiv\text{C}$
321	cyclo- $\text{C}_3\text{H}_5\text{OC}(\text{CH}_3)(\text{C}_2\text{H}_5)\text{C}\equiv\text{C}$
322	cyclo- $\text{C}_3\text{H}_5\text{OC}(\text{CH}_3)_2\text{C}\equiv\text{C}$
323	3- $\text{NO}_2\text{C}_6\text{H}_5$
324	4- $\text{NO}_2\text{C}_6\text{H}_5$
325	3- $(\text{CH}_3)_2\text{CHOC}_6\text{H}_5$

Table 14

Compound	R ¹
326	(CH ₃) ₂ C(CH ₃ O)C≡C
327	cyclo-C ₄ H ₇ C≡C
328	2-cyclopentyl-2-methyl-1-ethynyl
329	6-trifluoromethyl-2-pyridyl
330	6-phenoxy-2-pyridyl
331	6-benzyloxy-2-pyridyl
332	6-ethyl-2-pyridyl
333	6-propyl-2-pyridyl
334	6-isopropyl-2-pyridyl
335	6-butyl-2-pyridyl
336	6-isobutyl-2-pyridyl
337	6-sec-butyl-2-pyridyl
338	6-tert-butyl-2-pyridyl
339	2-cyclopentyl-2-ethyl-1-ethynyl
340	4-phenyl-2-pyrimidinyl
341	4-phenoxy-2-pyrimidinyl
342	4-benzyloxy-2-pyrimidinyl
343	4-methyl-2-pyrimidinyl
344	4-ethyl-2-pyrimidinyl
345	4-trifluoromethyl-2-pyrimidinyl
346	4-methoxy-2-pyrimidinyl
347	4-cyano-2-pyrimidinyl
348	4-propyl-2-pyrimidinyl
349	4-isopropyl-2-pyrimidinyl
350	4-sec-butyl-2-pyrimidinyl

Table 15

Compound	R ¹
351	4-butyl-2-pyrimidinyl
352	4-isobutyl-2-pyrimidinyl
353	4-tert-butyl-2-pyrimidinyl
354	4-pentyl-2-pyrimidinyl
355	4-hexyl-2-pyrimidinyl
356	4-cyclopropyl-2-pyrimidinyl
357	6-methyl-4-pyrimidinyl
358	6-ethyl-4-pyrimidinyl
359	6-propyl-4-pyrimidinyl
360	6-isopropyl-4-pyrimidinyl
361	6-butyl-4-pyrimidinyl
362	6-methoxy-4-pyrimidinyl
363	6-phenoxy-4-pyrimidinyl
364	6-benzyloxy-4-pyrimidinyl
365	6-ethoxy-4-pyrimidinyl
366	6-propoxy-4-pyrimidinyl
367	6-pentyl-2-pyridyl
368	6-hexyl-2-pyridyl
369	2-pyrazinyl
370	6-methyl-2-pyrazinyl
371	6-ethyl-2-pyrazinyl
372	6-propyl-2-pyrazinyl
373	6-butyl-2-pyrazinyl
374	6-phenoxy-2-pyrazinyl
375	6-benzyloxy-2-pyrazinyl

The compound of the invention of the formula [I-b] (Table 16 to 17)





Table 16

Compound	R ¹	R ^a
376	phenyl	F
377	phenyl	Cl
378	phenyl	CF ₃
379	4-chlorophenyl	F
380	4-chlorophenyl	Cl
381	4-chlorophenyl	CF ₃
382	(CH ₃) ₃ CC≡C	F
383	(CH ₃) ₃ CC≡C	Cl
384	(CH ₃) ₃ CC≡C	CF ₃
385	2-methylphenoxy	Cl
386	3-methylphenoxy	Cl
387	4-methylphenoxy	Cl
388	2-methylphenoxyethyl	Cl
389	3-methylphenoxyethyl	Cl
390	4-methylphenoxyethyl	Cl
391	2-methylbenzyloxy	Cl
392	3-methylbenzyloxy	Cl
393	4-methylbenzyloxy	Cl
394	2-pyrazinyl	Cl
395	2-pyrimidinyl	Cl
396	4-pyrimidinyl	Cl
397	1-pyrazolyl	Cl
398	CH ₃ ON=C(CH ₃)	Cl
399	C ₂ H ₅ ON=C(CH ₃)	Cl
400	6-benzyloxy-2-pyrazinyl	Cl

Table 17

Compound	R ²	R ^a
401	4-CH ₃ C ₆ H ₄ CH ₂ ON=C(CH ₃)	Cl
402	3-CH ₃ C ₆ H ₄ CH ₂ ON=C(CH ₃)	Cl
403	2-CH ₃ C ₆ H ₄ CH ₂ ON=C(CH ₃)	Cl
404	3,4-(CH ₃) ₂ C ₆ H ₄ CH ₂ ON=C(CH ₃)	Cl
405	2,5-(CH ₃) ₂ C ₆ H ₄ CH ₂ ON=C(CH ₃)	- Cl
406	6-trifluoromethyl-2-pyridyl	Cl
407	6-phenoxy-2-pyridyl	Cl
408	6-benzyloxy-2-pyridyl	Cl
409	6-ethyl-2-pyridyl	Cl
410	6-propyl-2-pyridyl	Cl
411	6-isopropyl-2-pyridyl	Cl
412	6-butyl-2-pyridyl	Cl
413	6-isobutyl-2-pyridyl	Cl
414	6-sec-butyl-2-pyridyl	Cl
415	6-tert-butyl-2-pyridyl	Cl
416	4-methyl-2-pyrimidinyl	Cl
417	4-ethyl-2-pyrimidinyl	Cl
418	4-trifluoromethyl-2-pyrimidinyl	Cl
419	4-methoxy-2-pyrimidinyl	Cl
420	6-methyl-2-pyrazinyl	Cl
421	6-ethyl-2-pyrazinyl	Cl
422	6-propyl-2-pyrazinyl	Cl
423	6-butyl-2-pyrazinyl	Cl
424	6-phenoxy-2-pyrazinyl	Cl
425	6-benzyloxy-2-pyrazinyl	Cl

The compound of the invention of formula [I-c] (Table 18)



5

Table 18

Compound	Y	E	G	Z
426	CH	CH	CH	N
427	CH	CH	N	CH
428	CH	N	CH	CH
429	N	CH	CH	CH
430	CCH ₃	CH	CH	N
431	CCH ₃	CH	N	CH
432	CCH ₃	N	CH	CH
433	CH	CH	N	N
434	CH	N	CH	N
435	N	CH	CH	N
436	N	CH	N	CH
437	CH	N	N	CH
438	CCH ₃	CH	N	N

Next, examples of the formulation are shown. In these examples, "part(s)" represents "part(s) by weight", the compounds of the invention are represented by Compound Numbers in Table 1 to 18.

Formulation Example 1

Fifty (50) parts of each of the Compounds 1 to 438, 3 parts of calcium ligninsulfonate, 2 parts of sodium laurylsulfate, and 45 parts of synthetic hydrated silica are well pulverized and mixed, to separately obtain wettable powders of each compound.

Formulation Example 2

Twenty (20) parts of each of the Compounds 1 to 438, 1.5 parts of sorbitan trioleate and 28.5 parts of an aqueous solution including 2 parts of polyvinyl alcohol are well mixed and wet pulverized. Then 40 parts of an aqueous solution including 0.05 part of xanthan gum and 0.1 part of aluminum magnesium silicate are added to the mixture, and 10 parts of propylene glycol are added, and the mixture is well mixed to give separately a flowable of each compound.

10 Formulation Example 3

Two (2) parts of each of the Compounds 1 to 438, 88 parts of kaolinite clay and 10 parts of talc are well pulverized and mixed to give separately a dust formulation of each compound.

Formulation Example 4

15 Five (5) parts of each of the Compounds 1 to 438, 14 parts of polyoxyethylenestyryl phenyl ether, 6 parts of calcium dodecylbenzenesulfonate, and 75 parts of xylene are well mixed to give separately an emulsifiable concentrates of each compound.

20 Formulation Example 5

Two (2) parts of each of the Compounds 1 to 438, 1 part of synthetic hydrated silicon oxide, 2 parts of calcium lignin sulfonate, 30 parts of bentonite, and 63.5 parts of kaolinite clay are well pulverized, and water is added. Then the mixture is well kneaded and granulated and dried to give separately a granule of each compound.

Formulation Example 6

Ten (10) parts of each of the Compounds 1 to 438, 35 parts of white carbons including 50% of ammonium polyoxyethylenealkylethersulfate salts and 55 parts of water

are well mixed and wet pulverized to give separately a flowable of each compound.

Formulation Example 7

5 Ten (10) parts of each of the Compounds 1 to 438, 5 parts of Sorpol 1200 (surfactant produced by Toho Chemical Co., Ltd.), 47 parts of dimethylsulfoxide and 38 parts of xylene are well mixed to give separately an emulsifiable concentrates of each compound.

10 Next, usefulness of the compounds of the invention as an active ingredient of an agricultural and horticultural fungicidal composition is shown by Test Examples.

Test Examples 1 to 6 are tests using a large amount (about 2000 L/ha) of spray on plants, and the extent of spraying is 15 so that the whole surface of a plant body is completely wet and the sprayed liquid drips.

Test Examples 7 and 8 are tests using a small amount (corresponding 200 L/ha) of spray on plants, in which fine small liquid drops of a spray are uniformly adhere to the surface 20 of a plant body and the extent of spraying is insufficient to for complete wetting of the surface of a plant body.

Controlling on a wheat disease in a field is so conducted in some cases that the spray liquid amount is about 200 L/ha. In this case, the amount of the sprayed liquid is insufficient 25 for wetting of the whole plant body, and the sprayed liquid is adheres on a plant as fine liquid drops. For exerting high control effect on the subject diseases even under such spray condition, suitable physicochemical properties such as permeation transfer property, volatility and the like are 30 required in addition to a basic effect.

In the following tests, compounds of the invention are represented by compound numbers described in Tables 1 to 8. The control effect of a compound of the invention was evaluated by visually observing the area of a lesion on a sample plant in investigation and comparing the area of a lesion in a non-treatment district and the area of a lesion in a districts treated with the compound of the invention.

Test Example 1: *Pyricularia oryzae* control test

Sand loam was compacted in a plastic pot, a rice (NIHONBARE) was seeded and grown in a green house for 20 days. Then, the Compounds 1, 3, 4, 6, 7, 9, 18, 19, 20, 24, 47, 104, 105, 113, 117, 118, 120, 294, 326, 329, 343 and 345 were formulated into flowable formulations according the Formulation Example 6, then, diluted with water to provide given concentration, and these were sprayed onto stems and leaves so as to give sufficient adhesion on the surface of rice leaves. After spraying, the plant was air-dried, and a suspension of *Pyricularia oryzae* was inoculated by spraying. After inoculation, the plant was left for 6 days at 28°C under high humidity, then, the control effect was checked.

As a result, the lesion areas on plants in the 500 ppm treatment districts and 200 ppm treatment districts using compound 1 of the invention were not more than 10% of the lesion area of a non-treatment district. The lesion areas on the plant in the 200 ppm treatment districts of the Compounds 3, 4, 6, 7, 9, 18, 19, 20, 24, 47, 104, 105, 113, 117, 118, 120, 294, 326, 329, 343 and 345 were not more than 10% of the lesion area of a non-treatment district.

Test Example 2: *Erysiphe graminis* control test

Sand loam was compacted in a plastic pot, a wheat (NORIN

No. 73) was seeded and grown in a green house for 10 days. Then, the Compounds 1, 3, 4, 6, 7, 9, 18, 19, 20, 24, 47, 104, 105, 113, 117, 118, 120, 277, 294, 326, 329, 343 and 345 were formulated into flowable formulations according the Formulation Example 5 6, then, diluted with water to provide given concentration, and these were sprayed onto stems and leaves so as to give sufficient adhesion on the surface of wheat leaves of wheat seedling carrying a spread second leaf. After spraying, the plant was air-dried, and one day after, *Erysiphe graminis* spores 10 were inoculated by falling. After inoculation, the plant was placed under illumination at 23°C for 7 days, then, the control effect was checked.

As a result, the lesion areas on plants in the 500 ppm treatment districts and 200 ppm treatment districts using 15 compound 1 of the invention were not more than 10% of the lesion area of a non-treatment district. The lesion areas on the plant in the 200 ppm treatment districts of the Compounds 3, 4, 6, 7, 9, 18, 19, 20, 24, 47, 104, 105, 113, 117, 118, 120, 277, 294, 326, 329, 343 and 345 were not more than 10% of the lesion 20 area of a non-treatment district.

Test Example 3: *Puccinia striiformis* control test

Sand loam was compacted in a plastic pot, a wheat (NORIN No. 73) was seeded and grown in a green house for 10 days. Then, the Compounds 1, 3, 4, 6, 7, 9, 18, 19, 20, 24, 47, 104, 105, 25 113, 117, 118, 120, 277, 294, 326, 329, 343 and 345 were formulated into flowable formulations according the Formulation Example 6, then, diluted with water to provide given concentration, and these were sprayed onto stems and leaves so as to give sufficient adhesion on the surface of wheat leaves. After 30 spraying, the plant was air-dried, and *Erysiphe graminis* spores

were inoculated. After inoculation, the plant was first left for 1 day at 23°C in dark high humidity atmosphere, then, further left for 6 days under illumination, then, the control effect was checked.

5 As a result, the lesion areas on plants in the 500 ppm treatment districts and 200 ppm treatment districts using compound 1 of the invention were not more than 10% of the lesion area of a non-treatment district. The lesion areas on the plant in the 200 ppm treatment districts of the Compounds 3, 4, 6,
10 7, 9, 18, 19, 20, 24, 47, 104, 105, 113, 117, 118, 120, 277, 294, 326, 329, 343 and 345 were not more than 10% of the lesion area of a non-treatment district.

Test Example 4: *Leptosphaeria nodorum* control test

Sand loam was compacted in a plastic pot, a wheat (NORIN
15 No. 73) was seeded and grown in a green house for 10 days. Then, the Compounds 1, 3, 4, 6, 7, 9, 18, 19, 20, 24, 47, 104, 105, 113, 118, 120, 277, 329, 343 and 345 were formulated into flowable formulations according the Formulation Example 6, then, diluted with water to provide given concentration, and these were sprayed
20 onto stems and leaves so as to give sufficient adhesion on the surface of wheat leaves. After spraying, the plant was air-dried, and a suspension of *Leptosphaeria nodorum* spores was inoculated by spraying. After inoculation, the plant was first left for 4 days at 15°C in dark high humidity atmosphere,
25 then, further left for 7 days under illumination, then, the control effect was checked.

As a result, the lesion areas on plants in the 500 ppm treatment districts and 200 ppm treatment districts using compound 1 of the invention were not more than 10% of the lesion
30 area of a non-treatment district. The lesion areas on the plant

in the 200 ppm treatment districts of the Compounds 3, 4, 6, 7, 9, 18, 19, 20, 24, 47, 104, 105, 113, 118, 120, 277, 329, 343 and 345 were not more than 10% of the lesion area of a non-treatment district.

5 Test Example 5: *Plasmopara viticola* control effect test

Sand loam was compacted in a plastic pot, a grape (Berry A) was seeded and grown in a green house for 40 days. Then, the Compounds 1, 3, 4, 6, 7, 9, 18, 19, 20, 24, 47, 104, 105, 113, 117, 118, 120, 277, 294, 326, 329, 343 and 345 were formulated
10 into flowable formulations according the Formulation Example 6, then, diluted with water to provide given concentration (200 ppm), and these were sprayed onto stems and leaves so as to give sufficient adhesion on the surface of wheat leaves. After spraying, the plant was air-dried, and a suspension of
15 zoosporangia of *Plasmopara viticola* was inoculated by spraying. After inoculation, the plant was first left for 1 day at 23 °C under high humidity, then, further left for 6 days at room temperature, then, the control effect was checked.

As a result, the lesion areas on plants in the treatment
20 districts using the Compounds 1, 3, 4, 6, 7, 9, 18, 19, 20, 24, 47, 104, 105, 113, 117, 118, 120, 277, 294, 326, 329, 343 and 345 were not more than 10% of the lesion area of a non-treatment district.

Test Example 6: *Sphaerotheca fuliginea* control effect test

25 Sand loam was compacted in a plastic pot, a cucumber (SAGAMI HANJIRO) was seeded and grown in a green house for 12 days. Then, the Compound 1 was formulated into flowable fomulation according the Formulation Example 6, then, diluted with water to provide given concentration (200 ppm), and this
30 was sprayed onto stems and leaves so as to give sufficient

adhesion on the surface of cucumber leaves. After spraying, the plant was air-dried, and *Sphaerotheca fuliginea* spores were inoculated. After inoculation, the plant was first left for 12 days at 23°C, then, the control effect was checked.

5 As a result, the lesion area on plants in the treatment district using the Compound 1 was not more than 10% of the lesion area of a non-treatment district.

Test Example 7: *Erysiphe graminis* control effect test

 Sand loam was compacted in a plastic pot, a wheat (NORIN
10 No. 73) was seeded and grown in a green house for 10 days. Then, the Compound 1 was formulated into emulsifiable formulation according the Formulation Example 7, then, diluted with water to provide given concentration (1000 ppm), and this was sprayed onto stems and leaves at a spraying liquid amount corresponding
15 to 200 L/ha over wheat seedling carrying a spread second leaf. After spraying, the plant was air-dried, and one day after, *Erysiphe graminis* spores were inoculated by falling. After inoculation, the plant was placed under illumination at 23 °C for 7 days, then, the control effect was checked.

20 As a result, no lesion was recognized at all in the treatment district using the Compound 1.

Test Example 8: *Puccinia striiformis* control test

 Sand loam was compacted in a plastic pot, a wheat (NORIN
25 No. 73) was seeded and grown in a green house for 10 days. Then, the Compound 1 was formulated into emulsifiable formulation according the Formulation Example 7, then, diluted with water to provide given concentration (1000 ppm), and this was sprayed onto stems and leaves at a spraying liquid amount corresponding to 200 L/ha over wheat seedling carrying a spread second leaf.
30 After spraying, the plant was air-dried, and *Puccinia*

striiformis spores were inoculated. After inoculation, the plant was placed in dark high humidity atmosphere at 23°C for 1 day, further, for 6 days under illumination, then, the control effect was checked.

- 5 As a result, no lesion was recognized at all in the treatment district using the Compound 1.

Industrial Applicability

- 10 The compound of the invention has a excellent fungicidal activity.